

1 **Multi-element signature of cuttlefish and its potential for the discrimination of**
2 **different geographical provenances and traceability**

3

4 *Maria Olga VARRÀ^a, Lenka HUSÁKOVÁ^{b*}, Jan PATOCKA^b, Sergio GHIDINI^a,*
5 *Emanuela ZANARDI^{a*}*

6

7 *^aDepartment of Food and Drug, University of Parma, Parma, Via del Taglio, 10, Parma 43126, Italy*

8 E-mail addresses: mariaolga.varra@studenti.unipr.it (M. O. Varrà); sergio.ghidini@unipr.it (S.
9 Ghidini).

10 *^bDepartment of Analytical Chemistry, Faculty of Chemical Technology, University of Pardubice,*
11 *Studentska 573 HB/D, Pardubice, CZ-532 10, Czech Republic*

12 E-mail address: jan.patocka@upce.cz (J. Patočka)

13

14 ***CORRESPONDING AUTHORS:**

15 **Lenka Husáková** (L. Husáková)

16 Department of Analytical Chemistry, Faculty of Chemical Technology, University of Pardubice
17 Studentská 573 HB/D, Pardubice, CZ-532 10 (Czech Republic)

18 Tel. +420 466 037 029

19 Fax: +420 466 037 068

20 E-mail address: lenka.husakova@upce.cz

21

22 **Emanuela Zanardi** (E. Zanardi)

23 Department of Food and Drug, University of Parma, Parma, Strada del Taglio, 10, Parma 43126, Italy

24 Tel. +39.0521.902.760

25 E-mail address: emanuela.zanardi@unipr.it

26 **ABSTRACT**

27 The measurement and analysis of fifty-two elements by quadrupole inductively coupled plasma mass
28 spectrometry (Q-ICP-MS) and direct mercury analysis were applied to origin discrimination of Italian
29 traditional cuttlefish (Chioggia, Venice lagoon) from Mediterranean and Atlantic samples. A total 68
30 specimens were analyzed in triplicates to generate 204 mass spectra profiles which were statistically
31 processed by different chemometric techniques. Loading weights from principal component analysis as input
32 for linear discriminant analysis (LW-LDA), stepwise-LDA (S-LDA) and variable influence of projection-
33 partial least square discriminant analysis (VIP-PLS-DA) were used to classify samples while retaining the
34 lowest possible number of key variables. VIP-PLS-DA was found to be the best variable selection-
35 discriminant tool combo since the selected Na-Co-B-K-Cd-V-U-Rb-Ni-Ba-Cu-As-Sr-Mn-Mo-Li-Ca-
36 Mg-Se-Bi-Cs-P-Y elemental pattern allowed **the samples to be classified** with 100% sensitivity, specificity
37 and accuracy.

38

39 **Abbreviations**

40 Certified reference materials, CRMs; correlation analysis, CA; discriminant function, DF; high energy
41 Helium mode, HE He; inductively coupled plasma mass spectrometry, ICP-MS; inductively coupled plasma
42 optical emission spectrometry ICP-OES; internal standard, ISTD; kinetic energy discrimination, KED; linear
43 discriminant analysis, LDA; leave-one-out-cross-validation, LOOCV; loading weights-based linear
44 discriminant analysis, LW-LDA; method detection limit of the method, MDL; method limit of
45 quantification, MLOQ; microwave digestion, MWD; partial last square-discriminant analysis, PLS-DA;
46 principal component, PC; principal component analysis, PCA; quadrupole inductively coupled plasma mass
47 spectrometry, Q-ICP-MS; rare earth elements, REEs; relative standard deviation, RSD; root mean square
48 error from cross-validation, RMSECV; root mean square error of estimation, RMSEE; root mean square
49 error of prediction, RMSEP; stepwise- linear discriminant analysis, S-LDA; variable influence on projection,
50 VIP; variable influence of projection-partial least square discriminant analysis, VIP-PLS-DA.

51

52

53 **Keywords:** ICP-MS; element profiling; fishery products; geographical origin; chemometrics.

54 **1. Introduction**

55 The common cuttlefish (*Sepia officinalis*, L., 1758) is a highly valuable fishery product whose distribution
56 extends from the Eastern Atlantic (from the Baltic and North Seas to South Africa coasts) up to the whole
57 Mediterranean Sea. Large scale fishing operations of common cuttlefish led to an average production of **21**
58 **825 000 kg in 2018**, which ensured the product's supply all over the world (especially in Italy, Spain, Japan,
59 and South Korea), while smaller-scale fishing is particularly profitable in some Mediterranean areas where it
60 has a high impact on local economies (FAO, 2020). This activity is particularly developed in the fishing
61 village of Chioggia (north-western Adriatic Sea in the Eastern Mediterranean Basin), whose *S. officinalis*
62 production reached more than **454 000 kg in 2019** (Clodia database, 2017).

63 Chioggia cuttlefish is well known to be of high quality thanks to the preservation of a close link with the
64 territory and the traditional processing methods by local fish plants, which **allow** the products to be officially
65 certified as Italian Traditional Agri-food Product (P.A.T.) (Ministerial Decree, G.U. n. 48, 2014). Foodstuffs
66 with specific geographical ties (especially when a quality mark has been recognized) are in turn rated more
67 highly in terms of quality by the consumer, mostly because of lower environmental impact and higher
68 perceived safety of regional products.

69 Alongside with the compulsory labelling requirement established by European legislation concerning the
70 provision of detailed information about the geographical origin of fish and seafood (European Parliament
71 and Council of the European Union, 2013), the protection and the promotion of traditional foods must
72 involve origin authenticity assessment both to ensure traceability and prevent commercial frauds (Ortea &
73 Gallardo, 2015).

74 Complex multi-disciplinary and cross-disciplinary approaches are usually required to verify the geographical
75 origin of fish and seafood since both environment and genetics can affect the final characteristics of the
76 products (Abbas et al., 2018). Nevertheless, considering the association between the concentrations of
77 elements in fish tissues and those in the surrounding aquatic environment, the determination of the multi-
78 element profile of fish and seafoods can be regarded as a valid analytical strategy to guarantee fish origin
79 authenticity. Several factors such as species, size, age, sex, and sexual maturity can directly affect the
80 elemental composition of aquatic animals, but food resources, climate, presence of contaminants, and many

81 water quality parameters (*pH*, dissolved oxygen, alkalinity) are those which differ to a greater extent between
82 countries (Li, Boyd, & Sun, 2016; Smith & Watts, 2009).

83 Several studies reported the possibility to discriminate fishery and seafood products as croaker (Chaguri et
84 al., 2015), sea cucumber (Kang et al., 2018), shrimps (Ortea & Gallardo, 2015), crabs (Luo et al., 2019), and
85 bivalve mollusks such as mussels (Costas-Rodríguez, Lavilla, & Bendicho, 2010) and clams (Iguchi, Isshiki,
86 Takashima, Yamashita, & Yamashita, 2014) using element composition. In these works, the chemical
87 characterization of samples was based on minor and/or trace elements (Costas-Rodríguez et al., 2010; Iguchi
88 et al., 2014), a combination of major, minor, and trace elements (Kang et al., 2018), or a combination of
89 major and trace elements plus stable isotopes analysis (Chaguri et al., 2015; Ortea et al., 2015; Luo et al.,
90 2019).

91 In the range of analytical methods applied to the determination of elements, inductively coupled plasma-
92 mass spectrometry (ICP-MS) combined with different chemometrics techniques holds a unique position by
93 virtue of speed, sensitivity, dynamic range, and elemental coverage (Drivelos & Georgiou, 2012; Danezis,
94 Tsagkaris, Camin, Brusica, & Georgiou, 2016). In this context, several supervised and unsupervised
95 chemometric tools such as, principal component analysis (PCA) (Chaguri et al., 2015; Kang et al., 2018;
96 Ortea et al., 2015), cluster analysis and *k*-means hierarchical clustering (Ortea et al., 2015), linear
97 discriminant analysis (LDA) (Costas-Rodríguez et al., 2010; Iguchi et al., 2014; Ortea et al., 2015; Kang et
98 al., 2018; Luo et al., 2019), support-vector machine (Luo et al., 2019), soft-independent modelling of class
99 analogy and artificial neural networks (Costas-Rodríguez et al., 2010) were applied for authenticity and
100 traceability purposes.

101 The determination of elemental composition of fish and seafood to trace their geographical provenance is
102 particularly suitable for cephalopods mollusks. Cephalopods such as cuttlefish are in fact at high trophic
103 levels in the aquatic food chain and the overall elements accumulated in tissues are good indicators of the
104 surrounding habitat (Bosch, O'Neill, Sigge, Kerwath, & Hoffman, 2015). In addition, cephalopods are non-
105 migratory animals and, therefore, their elemental composition can reasonably be expected to be constant
106 during life (Gopi et al., 2019). Nevertheless, very few studies addressed the topic of origin discrimination of
107 cephalopods mollusks using elemental composition, in spite of being focused on the analysis of a limited
108 number of major and trace elements included in hard structures such as statoliths (Arbuckle & Wormuth,

109 2014) or ink (Bua et al., 2017). These components are in fact not always retained in the final commercial
110 product and, therefore, not always exploitable in practical food surveillance operations to trace back the
111 origin of cephalopods.

112 Based on this background, the present work aimed at outlining for the first time the *S. officinalis* multi-
113 elemental profile of edible tissues to verify whether useful information could be extracted and specifically
114 linked to the geographical origin of the Italian traditional cuttlefish from Chioggia (FAO subarea 37.2.1) in
115 order to be differentiated from non-Chioggia cuttlefish caught in other fishing areas of the Mediterranean Sea
116 (FAO 37.1/37.2) and in the French Atlantic Ocean (FAO 27.7.e). For this purpose, fifty-one elements were
117 determined by ICP-MS and Hg was quantified via atomic absorption spectroscopy. Unsupervised and
118 supervised pattern recognition methods were used to investigate sample characteristics, identify the key
119 discriminant elements, and, at the same time, develop classification rules.

120 **2. Materials and Methods**

121 *2.1. Reagents and standards*

122 The ultrapure water (0.055 $\mu\text{S cm}^{-1}$ conductivity) obtained using the Milli-Q[®] water purification system
123 (Millipore, Bedford, USA) was used for the preparation of all solutions. Sub-boiled nitric acid was prepared
124 from nitric acid (65%, w/w) of Selectipur quality (Lach-Ner, Neratovice, Czech Republic) using the
125 distillation equipment BSB-939-IR (Berghof, Eningen, Germany). Hydrogen peroxide (Trace Select, $\geq 30\%$,
126 w/w) was purchased from Fluka Chemie AG (Buchs, Switzerland).

127 Multi-element stock solution “A” containing 10 mg L⁻¹ of Li, B, Al, V, Cr, Fe, Ni, Co, As, Se, Rb, Sr, Zr,
128 Mo, Ru, Pd, Cd, Sn, Sb, Cs, Ba, Hf, Re, Pt, Tl, Pb, Bi, and Th was prepared from the Supelco ICP multi-
129 element standard solution IV (Merck, Darmstadt, Germany) and single element standards of concentration
130 $1 \pm 0.002 \text{ g L}^{-1}$ (Analytika Ltd., Prague, Czech Republic or SCP Science, Montreal, Canada). Multi-element
131 solution “B” containing 1 mg L⁻¹ of La, Ce, Pr, Nd, U (“B1”) and 0.20 mg L⁻¹ of Y, Tb, Ho, Yb, Sm, Eu, Gd,
132 Er, Lu, and Dy (“B2”) was prepared from the stock solution of rare earth elements Astatol mix “M008”
133 (Analytika Ltd., Prague, Czech Republic). Multi-element solution “C” containing 50 mg L⁻¹ of Na, Mg, P, K,
134 Ca, Mn, Cu, and Zn was prepared from single element standards of 1 g L⁻¹ obtained from Analytika Ltd. The

135 internal standard solution (ISTD) was prepared from 1 g L⁻¹ stock solution of Rh obtained from SCP Science
136 (Montreal, Canada). Carbon reference solutions were prepared from 10 g L⁻¹ of C stock solution prepared
137 from urea of TraceSelect quality (Fluka Chemie AG, Buchs, Switzerland).

138 2.2. Sampling and preparation of cuttlefish

139 A total of 68 samples of common cuttlefish (*S. officinalis*, L.) including n = 17 samples from Adriatic Sea
140 (Chioggia, Italy, FAO 37.2.1), n = 25 non-Chioggia samples from the Mediterranean Basin (FAO 37.1/37.2),
141 and n = 26 samples from North-eastern Atlantic Ocean (France, FAO 27.7.e) were analyzed. Specimens were
142 caught by fishing trawlers during the months of September and randomly collected from different batches in
143 local fish plants located in Chioggia (Venice, Italy), by choosing homogeneous sizes and weights (mantle
144 lengths from 10 ± 2 cm; body weight 125 ± 25 g). After collection, samples were immediately frozen and
145 stored at -20 °C for around 3 months prior to be further processed for multi-element analysis.

146 2.3. Quality assurance and quality control

147 The following commercially supplied certified reference materials (CRMs) were analyzed: NIST SRM 1577
148 Bovine Liver (National Institute of Science and Technology, NIST, Gaithersburg, MD, USA); NIST SRM
149 1566 Oyster Tissue (NIST, Gaithersburg, MD, USA); BCR[®] certified reference material (CRM)184 Bovine
150 muscle (Institute for Reference Materials and Measurements, IRMM, Geel, Belgium); BCR[®] 185 Bovine
151 Liver (IRMM, Geel, Belgium); CRM NCS ZC73015 Milk Powder (National Research Centre for Certified
152 Reference Materials, NRCRM, Beijing, China); P-WBF CRM 12-2-04 Essential and Toxic Elements in
153 Wheat Bread Flour (pb-anal, Kosice, Slovakia); CRM12-2-03 P-Alfalfa Essential and toxic elements in
154 Lucerne (pb-anal, Kosice, Slovakia); SMU CRM 12-02-01 Bovine liver (pb-anal, Kosice, Slovakia).

155 2.4. Sample preparation

156 2.4.1. Pre-processing of cuttlefish

157 Frozen samples were thawed overnight (16–18 h at +4 °C) before processing for multi-element analysis.
158 Each specimen was then washed with deionized water and skin, cuttlebone, gills, reproductive and digestive

159 tracts were carefully removed without causing their rupture. The head, arms and tentacles were excluded,
160 while the mantle and the lateral fins were rinsed again with ultrapure water and minced with a ceramic knife.

161 2.4.2. Freeze-drying process

162 Pre-treated **sample** portions of approx. 10 g were individually transferred into 100 mL cleaned round bottom
163 flasks wherein the material was dried and placed into a deep freezer at $-80\text{ }^{\circ}\text{C}$ for 24 hours to provide a
164 necessary conditioning for drying. Lyophilization was conducted at $-50\text{ }^{\circ}\text{C}$ and 0.01 bar for 24 h with a LIO-
165 5P apparatus (CinquePascal srl, Trezzano sul Naviglio, Milan, Italy). Dried samples were subsequently
166 removed from the flasks, homogeneously **ground** into a powder by using ceramic mortars and pestles and
167 then individually sealed into LDPE bags.

168 2.4.3. Microwave assisted digestion

169 Microwave digestion (MWD) of samples was carried out by using the SpeedwaveTM MWS-3⁺ (Berghof,
170 Eningen, Germany) microwave system with the maximum total output of the microwave generator (1450 W)
171 and equipped with the optical sensor technology for contactless real-time recording of the sample
172 temperature. The high-pressure resistant (up to 100 bar) PTFE vessels DAC-100S, together with the
173 Multitube System (all from Berghof, Eningen, Germany), were used for sample digestion. This arrangement
174 allowed the simultaneous digestion of three samples in one DAC-100S PTFE vessel by placing three PFA
175 tubes into each vessel (Husáková et al., 2015). The maximal number of used DAC-100S vessels for one
176 digestion cycle was eight.

177 A powdered cuttlefish or CRM sample aliquot of 0.1 g was weighed into a 10 mL PFA tube. Then, 4 mL of
178 16% HNO_3 (65%, w/w HNO_3 , 1:3 diluted) and 1 mL of 30% H_2O_2 were added, leaving the vessels open until
179 the initial reaction subsided. Three PFA tubes containing the same sample and reagents were placed into the
180 outer 100mL PTFE digestion vessel previously filled with 25 mL of HNO_3 (16%, v/v), by ensuring that the
181 level of liquid in the outer PTFE vessel was higher than those in the PFA tubes. This way, the vapor
182 pressures were compensated and the evaporation of the solution from the PFA tubes was avoided (Husáková
183 et al., 2015). The samples were digested following a five-step program: (i) 20 min at $180\text{ }^{\circ}\text{C}$ and 80 % power
184 (ramp 5 min), (ii) 20 min at $220\text{ }^{\circ}\text{C}$ and 95 % power (ramp 5 min), (iii–v) 5 min at $100\text{ }^{\circ}\text{C}$ and 10 % power
185 (ramp 1 min). The resulting colorless solutions were diluted to 25 mL with deionized water.

186 Each sample was mineralized in three replicates. Blanks, consisting of deionized water and reagents were
187 subjected to a similar preparation procedure.

188 To assess the properness of the digestion process, the residual carbon content in the final cuttlefish digests
189 was determined by a previously described inductively coupled plasma optical emission spectrometry (ICP-
190 OES) method (Husáková et al., 2011) was 5.5 ± 0.4 % (n = 3).

191 *2.5. ICP multi elemental analysis*

192 ICP-MS measurements were performed by using the Agilent 7900 quadrupole mass spectrometer (Q-ICP-
193 MS, Agilent Technologies, Inc., Santa Clara, CA, USA) equipped with an octopole-based collision cell for
194 interference removal using kinetic energy discrimination (KED).

195 The standard sample introduction system, consisting of a glass concentric nebulizer MicroMist ($400 \mu\text{L min}^{-1}$)
196 ¹), the Peltier-cooled (2°C) Scott quartz spray chamber and quartz torch with 2.5 mm internal diameter
197 injector, was used. For precise delivery of samples and ISTD, a low-pulsation, 10-roller peristaltic pump
198 with three separate channels was employed. The internal standard kit, including connecting tubing,
199 connectors and the “Y” piece was used for simultaneous internal standard aspiration and its mixing with the
200 sample. The standard sampling and skimmer nickel cones with orifices of 1 and 0.45 mm, respectively, were
201 used. ICP-MS operating conditions were optimized during each start-up sequence by using the multi-
202 elemental tuning solution (Agilent Technologies, Inc., Santa Clara, CA, USA) containing $1 \mu\text{g L}^{-1}$ of Ce, Co,
203 Li, Mg, Tl and Y, in order to obtain the highest possible sensitivity for elements of low, middle and high m/z.
204 Using the typical operating conditions summarized in Table 1, a sensitivity of $6000 \text{ counts s}^{-1} \text{ per } \mu\text{g L}^{-1}$ and
205 a resolution of 0.64 amu peak width (full width at half maximum intensity) were achieved for $^7\text{Li}^+$. The same
206 parameters were $50000 \text{ counts s}^{-1} \text{ per } \mu\text{g L}^{-1}$ and 0.62 for $^{89}\text{Y}^+$, and $30000 \text{ counts s}^{-1} \text{ per } \mu\text{g L}^{-1}$ and 0.60 for
207 $^{205}\text{Tl}^+$.

208 For sample analysis the “General Purpose“ plasma mode included in the ICP-MS MassHunter software was
209 used (Agilent Technologies, Inc., Santa Clara, CA, USA). The working parameters of the cell mode “no-gas”
210 were autotuned during the instrument start-up sequence. The working parameters of the collision cell for
211 helium (“He”) and high energy He (“HE He”) modes were adjusted manually. The time required for a

212 transition between cell modes was 5 s. Parameters related to sample introduction and plasma conditions were
213 consistent for all modes (see Table 1).

214 Concentrations of a total of 51 elements were evaluated from calibration curves with coefficients of
215 determination better than 0.999 and built up within the ranges given below.

216 Calibration solutions: blank, 1, 5, 10, 50, 100 $\mu\text{g L}^{-1}$ of Li, B, Al, V, Cr, Fe, Ni, Co, As, Se, Rb, Sr, Zr, Mo,
217 Ru, Pd, Cd, Sn, Sb, Cs, Ba, Hf, Re, Pt, Tl, Pb, Bi, Th; 0.1, 0.5, 1, 5, 10 $\mu\text{g L}^{-1}$ La, Ce, Pr, Nd, U; 0.02, 0.1,
218 0.2, 1, 2 $\mu\text{g L}^{-1}$ of Y, Tb, Ho, Yb, Sm, Eu, Gd, Er, Lu, and Dy; 0.5, 1, 5, 10 mg L^{-1} of Na, Mg, P, K, Ca, Mn,
219 Cu, and Zn. The solutions were prepared daily by appropriate dilution of multi-element solutions “A”
220 ($500 \mu\text{g L}^{-1}$), “B” ($50 + 10 \mu\text{g L}^{-1}$) and “C” (50mg L^{-1}) in 25 mL volumetric flasks (see Section 2.1).

221 To compensate possible instrumental drift and matrix effects, a $200 \mu\text{g L}^{-1}$ Rh ISTD was simultaneously
222 aspirated and mixed with samples.

223 2.6. Mercury analysis

224 For the quantification of Hg content, the direct solid sampling analysis using a single-purpose atomic
225 absorption spectrometer AMA 254 (Altec Ltd., Prague, Czech Republic), based on in situ dry ashing
226 followed by gold amalgamation atomic absorption spectroscopy, was used. Each time, samples were
227 weighed in a nickel boat and analyzed under the following experimental conditions: typical sample mass,
228 50 mg; drying time, 60 s; decomposition, $\sim 750 \text{ }^\circ\text{C}$ for 150 s; waiting step, $900 \text{ }^\circ\text{C}$, 45 s, necessary for
229 quantitative release of trapped mercury from the gold amalgamator to the measuring cuvette. The peak area
230 absorbance at 253.6 nm was monitored. The flow rate of oxygen (99.5%) carrier gas was 170 mL min^{-1} .

231 The AMA 254 spectrometer was regularly calibrated using blank and four working standard solutions
232 containing Hg(II) ranging from 0.025 to 0.2 mg L^{-1} . The working standard solutions were prepared from
233 standard aliquots of 10 mg L^{-1} prepared from a 1 g L^{-1} stock solution of Hg(II) (Analytika Ltd., Prague,
234 Czech Republic).

235 2.7. Data treatment

236 Statistic was applied to elemental concentrations referring to cuttlefish dry matter (d.m.) and carried out
237 using IBM-SPSS (v. 23.0, SPSS Inc., Chicago, IL, USA) SIMCA-P v.14.1 (Umetrics, Umeå, Sweden), and
238 Statgraphics Centurion v.18.1 (StatPoint Technologies, Inc, Warrenton, VA, USA) software. The normality

239 and homogeneity of variance of data was **checked beforehand** by applying the Shapiro-Wilk's and Levene's
240 tests, respectively ($p \leq 0.05$). Since the normal distribution assumption was violated for most of the
241 elemental data, quantitative differences between groups of cuttlefish from different origins were investigated
242 by the nonparametric Kruskal–Wallis test with Dunn's multiple post hoc test for multiple comparison ($p \leq$
243 0.05). Data **was** presented as mean values \pm margin of error (calculated at 95% confidence level), median and
244 minimum and maximum values found.

245 While **the nonparametric statistic** was used to compensate for the lack of data normal distribution, thus
246 preserving the original information and avoiding the loose of any intuitive meanings when describing
247 element concentrations resulting from univariate data analysis, a Box-Cox transformation was instead
248 applied to normalize data for subsequent bivariate and multivariate data analyses.

249 First, the bivariate Pearson's correlation analysis (CA) was employed to Box-Cox transformed data to
250 explore variable distribution and look for strong positive ($r > 0.6$, $p \leq 0.01$) or strong negative ($r < -0.6$, $p \leq$
251 0.01) linear correlations between binary variables.

252 Because of **the** highly different magnitude of values among element concentrations, the Box-Cox data were
253 then scaled applying a Z-score standardization and further processed by multivariate data analyses. PCA was
254 used for further investigation of covariance patterns in elemental concentrations, data's first interpretation
255 and multivariate outliers' removal based on Hotelling's T^2 test results (95% confidence interval). During this
256 step, the number of variables was also reduced by selecting elements characterized by loading values
257 (rescaled to $0-1$) > 0.70 .

258 Afterwards, supervised classifiers combined with different variable selection procedures were developed to
259 achieve the discrimination of samples according to the geographical origin by using the lowest number and
260 the most influential combination of variables. To that end, linear discrimination based on variable loading
261 weights previously selected by PCA (LW-LDA), forward stepwise-LDA (S-LDA) and variable influence on
262 projection-(VIP)-PLS-DA were independently tested and compared in terms of prediction ability. All the
263 models were cross-validated using leave-one-out-cross-validation (LOOCV) to exclude overfitting. In
264 addition to LOOCV, the validity of the PLS discriminant models was also checked by permutation testing
265 (400 random permutations).

266 When supervised modelling was performed, about 33% of data (n = 68) was randomly sidelined from the
267 calibration set (n = 136) and used for independent external validation. Models' overall performances in
268 internal and external validation were evaluated in terms of accuracy (%), sensitivity (%), and specificity (%).
269 Further details about the statistical treatments applied can be found in Supplementary material,
270 Supplementary Experimental Section.

271 **3. Results**

272 *3.1. Analytical performances and validation*

273 The analytical determination of fifty-two major, trace and ultra-trace elements in mineralized cephalopod
274 samples was carried out by Q-ICP-MS, taking advantage of its well-known sensitivity toward the targeted
275 elements. On the other hand, analysis of Hg by thermal decomposition was performed by AMA254 without
276 pre-treatment and/or preconcentration steps, thus overcoming the most serious problems related to ICP-MS
277 analysis of Hg, e.g. long washout time, non-linear calibration curves, and decreasing sensitivity with time (Li
278 et al., 2006).

279 The microwave-assisted pressurized digestion which involved small sample amounts (100 mg) was
280 employed to ensure rapid sample preparation for subsequent ICP-MS analysis while attaining the high
281 decomposition efficiency. One of the most important aspects regarding the employed MWD method is the
282 possibility to use diluted solutions for sample preparation, thus reducing the quantity of reagents, risks of
283 contamination and generation of residues. These are important parameters for the development of greener
284 analytical methods (Gonzalez et al., 2009). Moreover, MWD plays an important role in decreasing some
285 kinds of spectral and non-spectral interferences, such as those deriving from carbon containing species or
286 different chlorides, by conversion to significantly less or non-interfering species (Husáková et al., 2011;
287 Bizzi et al., 2017). He mode with KED or HE He mode with higher cell gas pressure and higher energy
288 discrimination voltages, were used for the analytes that suffered from spectral effects (i.e. V, Cr, Mn, Co, Ni,
289 Fe, Co, Cu, As, and Se) caused by polyatomic ions deriving from the plasma gas, sample solvent and other
290 sample matrix elements such as Na, K, Ca, Mg, P, S, C, and Cl (see Table 1 and Supplementary Materials,
291 **Table S1**). He and/or HE He cell modes had the additional benefit of reducing the response for low mass

292 matrix elements like Na, K, Ca or P by an order of magnitude ($^{23}\text{Na}^+$, $^{44}\text{Ca}^+$) or more ($^{39}\text{K}^+$, $^{31}\text{P}^+$), thus
293 effectively raising the upper linear range for these elements. Thereby elements that would have needed to be
294 analyzed by ICP-OES, were included in the ICP-MS.

295 As for the ISTD, Rh was chosen for its mid-range mass and ionization potential and because of its absence in
296 the analyzed samples. Recoveries of Rh, relative to the initial calibration blank for the entire 8-hour sequence
297 of sample digests, are shown in Supplementary Materials, Fig. S1. ISTD behavior across the mass range was
298 found to be very consistent over time and downward drift was considered acceptable.

299 Data accuracy was evaluated by the analysis of different CRMs (NIST SRM 1577 Bovine Liver, NIST 1566
300 Oyster Tissue, BCR CRM 184 Bovine muscle, BCR CRM 185 Bovine Liver, CRM 12-2-01 Bovine Liver),
301 primarily intended for the evaluation of analytical methods and instruments used for the determination of the
302 mass fraction values of selected elements in marine tissue, foods, or similar materials. Since the levels of
303 most lanthanides and actinides are not certified in these CRMs, three additional certified standards (NCS ZC
304 73015 Milk Powder, CRM 12-2-04 Wheat bread flour, and CRM 12-2-03 Essential and toxic elements in
305 Lucerne) were analyzed to validate data for these elements. The high level of agreement between target and
306 found values demonstrated trueness of data obtained (Supplementary Materials, Table S2 and Fig. S3).

307 The precision of the method was evaluated in terms of intra-day and inter-day comparison. Intra-day
308 precision was determined by analysis of individual control materials three times during the same day. Inter-
309 day precision was determined by analysis of the same standards on three different days over a period of one
310 month. Within each series, every solution was analyzed in triplicate. Relative standard deviation (RSD) was
311 calculated for both series of analyses. The RSD values of intra-day and inter-day studies were mostly found
312 to be below 14% thus showing a good precision of the method (Supplementary Materials, Table S2).

313 Method detection limits (MDLs) and method quantification limits (MLOQs) - reported in Supplementary
314 Material, Table S3 - were evaluated as a triple and tenfold standard deviation, respectively, of ten
315 consecutive measurements of blank signal divided by the slope of the calibration curve. Detection limits
316 were found to be low enough that selected elements could be determined at the background level. Fig. S2
317 and Table S3 (Supplementary Materials) also summarize relative sensitivities of Q-ICP-MS for analysis of
318 individual elements with the use of Rh ISTD.

319 *3.2. Concentrations of elements in cuttlefish samples*

320 The concentrations of elements measured in the present study were presented and discussed as median values
321 considering the non-normal distribution of the raw data. Results from descriptive statistics are listed in Table
322 2. The most abundant elements were found to be Na, Mg, P and K, whose concentration levels were higher
323 than 1% of weight and quite variable among cuttlefish from different countries. According to results from
324 Kruskal–Wallis test, Na and Mg amounts were significantly higher ($p \leq 0.05$) in samples from FAO 27.7.e
325 (French Atlantic) compared to samples from FAO 37.1/37.2 (Mediterranean area) and FAO 37.2.1
326 (Chioggia). At the same time, an opposite trend was observed for P and K, for which the highest
327 concentrations ($p \leq 0.05$) were found in Chioggia cuttlefish (Table 2).

328 Concentrations between 1 and 0.01% of weight (10000–100 mg kg⁻¹, d.m.) were found for two elements
329 corresponding to Li and Ca, which showed significantly higher median concentrations in French Atlantic
330 samples (0.51 mg kg⁻¹ and 1760 mg kg⁻¹, respectively) ($p \leq 0.05$) compared to the other groups. The wide
331 variability in both major and minor element contents found in cuttlefish specimens of different geographical
332 origins and within samples of the same origin can be attributed to their natural variation in seawater, but the
333 assimilation by marine animals of Na, Mg, and Ca from the organic matter in the aquatic environment also
334 varies with the feeding status and the age of the animal (Lall, 2002). Moreover, the presence of P in fish and
335 seafood tissues can be directly attributable to the food sources since its concentration in seawater is lower
336 compared to the other elements (Lall, 2002).

337 The main trace elements, ranging between 100 and 1 mg kg⁻¹, followed the decreasing order Zn > Al > Sr >
338 Cu > Fe > B > Rb > Se > Mn in samples from Chioggia, Zn > Cu > Sr > Fe > Al > B > Rb > Mn > Se in
339 samples from the Mediterranean area, and Zn > Sr > Cu > B > Fe > Al > Rb > Se > Mn in samples from the
340 French Atlantic. Among these elements, only Se, Rb, and Sr were different in median concentrations
341 according to all three geographical regions ($p \leq 0.05$) and the concentration ranges were in line with those
342 previously published in literature for *S. Officinalis* (Raimundo, Pereira, Vale, & Caetano, 2005; Ayas &
343 Ozogul, 2011).

344 Different median contents varying between 62.3 and 75.8 mg kg⁻¹ were also found for As ($p \leq 0.05$). The
345 wide presence of arsenic in the aquatic environment is linked both to the geochemical characteristics of the
346 region and anthropogenic activities, especially those related to agricultural practices (Neff, 2002). Due to
347 toxic properties, some concerns regarding the dietary exposure to arsenic have arisen in recent times, but

348 maximum levels for this metal in fishery products have not been established yet by European legislations.
349 Fish and seafood, in particular, have been reported as the largest food contributors to overall total arsenic
350 exposure, despite the largest proportion is represented by organic arsenic species which are known to be less
351 or no toxic compared to relative inorganic species (European Food Safety Authority, EFSA, 2009). In
352 addition, inorganic arsenic concentrations decrease with increasing content of total arsenic (European Food
353 Safety Authority, EFSA, 2009).

354 Most of the elements analyzed in the present paper were found to be present in cuttlefish tissues at mean
355 concentration lower than 1 mg kg⁻¹. Levels of ultra-trace elements as rare earth elements (REEs) were lower
356 than 100 µg kg⁻¹ on average, with the highest median concentration for Ce (24.0 µg kg⁻¹) and the lowest one
357 for Lu (0.047 µg kg⁻¹) both in in Mediterranean samples (Table 2). The natural pattern of REEs in the
358 aquatic environment, which is associated to the mobilization of nutrients from the underlying soils, can be
359 permanently altered by some anthropic activities and effectively used to investigate the geographical
360 provenance of marine animals (Noack, Dzombak, & Karamalidis, 2014). In the present work, the
361 significantly different concentrations of Tb, Dy, Ho, Er, and Yb geographically discriminated Chioggia from
362 French specimens, while those of La, Pr, Nd, Eu, and Gd discriminated Mediterranean from French
363 specimens ($p \leq 0.05$). In accordance with these results, the amount of both La and Ho was previously
364 reported to be potentially useful to authenticate fish samples from different areas in the Mediterranean Sea
365 (Varrà, Ghidini, Zanardi, Badiani, & Ianieri, 2019).

366 Regardless the origin, all the cuttlefish samples analyzed in the present work were in line with the maximum
367 limits for toxic heavy metals established by European regulations No. 1881/2006/EC and subsequent
368 amendments and additions (Commission of the European Communities, 2006). Both cadmium and lead did
369 not exceed the threshold value of 1 mg kg⁻¹ set for the edible part of cephalopods, as well as Hg was always
370 found to be below the threshold limit of 0.5 mg kg⁻¹. The sample groups under investigation did not show
371 significant differences in Pb and Hg concentrations ($p > 0.05$), but Cd median amount in French Atlantic
372 specimens was found to be approximately 7 and 20 times higher than in Mediterranean and Chioggia
373 samples, respectively (see Table 2). Furthermore, the comparison with results reported by other authors for
374 cuttlefish from North eastern Atlantic and Adriatic Sea, showed that Hg, Pb, and Cd concentrations found in
375 the present work fell within the same ranges (Bustamante, Lahaye, Durnez, Churlaud, & Caurant, 2006;

376 Storelli, Giacomini-Stuffer, Storelli, & Marcotrigiano, 2007). Even in the case of heavy toxic metals, a
377 close link between their concentrations in seawater and environmental pollution of specific areas exists
378 (Carvalho, Santiago, & Nunes, 2005). This, together with Ni, Co, and Bi amounts, which also varied
379 significantly among sample groups, makes suggestions for future insights about the effective role and
380 contribution of these elements for the geographical origin assessment of fish and seafood.

381 *3.3. Chemometric classification of the geographical origin of cuttlefish*

382 Since no specific information concerning the best arrangement of elements to discriminate the geographical
383 origin of cephalopods is available in literature, the first stages of multivariate analysis took into consideration
384 all the fifty-two elements measured, starting with the data **exploration** by means of CA and PCA.
385 Nevertheless, multicollinearity between variables as well as geographical-unrelated variability are a source
386 of increasing noise, thus the reduction of the variables to the most predictive ones was subsequently
387 preferred, especially in view of a future practical implementation of the methodology.

388 The selection of variables is strictly dependent on the sample matrix properties and dataset overall
389 characteristics; hence its application needs to be evaluated on an individual basis. Although being the most
390 frequently used classification technique in studies dealing with element profiling, S-LDA is
391 coming under growing criticism because of the deceiving results deriving from the random rather than
392 effective significance of the variables extracted, especially when the number of potential explanatory
393 variables is high (Smith, 2018). On the other hand, one of the main problems related to the use of
394 multivariate data analysis in food authentication studies is just the difficulty in comparing the obtained
395 results with those already published but using a wide variety of different chemometrics techniques.
396 Therefore, in the context of the present work, S-LDA and LW-LDA, were used for classification by variable
397 selection, but outputs were compared with those obtained through the application of VIP-based PLS-DA.
398 The latter, in fact, consists of a more robust and flexible algorithm, particularly suited for the classification of
399 a large number of samples and is suggested to be a more powerful tool for reliable variable selection
400 compared to the traditional LDA (Rashid, Hussain, Ahmad, & Abdullah, 2019).

401 *3.3.1. Correlation and principal component analysis*

402 Pearson's associations between all the elements were reported by plotting the correlation matrix shown in
403 Fig. 1. The main significant patterns of covariations were found among most of the REEs and, in particular,
404 between Ce and La ($r > 0.9$), and among Nd, Pr, Tb, Ho, Gd, Er, Eu, Dy, and Sm ($r > 0.8$). Similarly,
405 bivariate positive correlation regarding the patterns Na–Sr–Cd–Li–Mg–Ca ($r > 0.9$), P–K ($r > 0.9$), and Cs–
406 Rb ($r > 0.8$) were also detected. Concentrations of U were positively correlated with Ca, Mg, Pd, Li, Cd, Sr,
407 and Na ($r > 0.8$) as well as amounts of Co with Cu, Mo, and V ($r > 0.7$). The most important and significant
408 negative correlations were instead found among K–Rb–Cs–Mn and among Na–Sr–Cd–Li–Ca–Mg–U ($r < -$
409 0.7) patterns.

410 The association between the characteristic element distributions and cuttlefish origins was further
411 investigated by PCA. The first PCA computation took into consideration all the fifty-two elements and, as a
412 result, the number of original variables was set by LOOCV to six final principal components (PCs) enclosing
413 72.6% and 57.3% of the explained and predictive data variance, respectively. The first two PCs of the model
414 contributed for 27.5% and 20.8% of the total explained variance, with the PC2 enclosing the fraction of
415 information related to sample provenance, as reported in Fig. 2A. The distribution of Chioggia cuttlefish
416 along the negative axis of the PC2 was mainly dominated by the highest contribution of Rb and K, followed
417 by Mn, Al, Cs, P, As, and, at lower levels, Bi (see loading vectors reported in Fig. 2A). On the other hand,
418 French Atlantic cuttlefish distribution was mainly guided by contribution from Na, Sr, Cd, Li, Mg, Ca, U,
419 and Mo, whose vectors angles nearness underlined the strong covariation previously stated by CA. Likewise,
420 a strong contribution and a high degree of covariation was confirmed in PCA for REEs, which impacted on
421 the negative axis of **PC1**.

422 During this preliminary stage, one sample from Chioggia was also found to be a strong outlier since it fell
423 outside the 95% confidence interval defined by the Hotelling's T^2 range (see Fig. 2A). For this reason, it was
424 excluded from subsequent statistics.

425 Since the most influential loading values were found for a total of 16 out 52 elements (Li, B, Na, Mg, Ca,
426 Co, Cu, Zn, Sr, Mo, Ru, Pd, Cd, Hf, Pt, and U), variable filtering was performed (see Section 2.7) and PCA
427 was thus recomputed. Loading weights (**0-1 scaled**) of selected elements are reported in Supplementary
428 Materials, Table S4. The reduced PCA model extracted three PCs that explained 84.5% of the data
429 variability. The PC1 (explained variability, 54.1%; predictive variability, 48.3%) was mainly characterized

430 by Zn, Cd, Pd, Ca, U, Sr, while in the PC2 (explained variability, 16.3%; predictive variability, 17.1%) Pt
431 was the most influential element (Fig. 2B). Hence, maximum variation in the original dataset was retained
432 while reducing the number of significant features. At the same time, sample separation in the score space
433 was significantly expressed along the PC1 using the selected variables rather than the whole set of elements.

434 3.3.2. Classification of cuttlefish by origin: LDA and PLS-DA approaches using selected variables

435 The LW-LDA classification model for cuttlefish origins, created by using variables previously selected by
436 PCA (Li, B, Na, Mg, Ca, Co, Cu, Zn, Sr, Mo, Ru, Pd, Cd, Hf, Pt, and U) was described by two discriminant
437 functions (DFs) explaining 99.7% of the data variability. The DF1 (explained variability, 83.1%; canonical
438 correlation, 97.2%) and the DF 2 (explained variability, 16.6%; canonical correlation, 86.7%), whose
439 statistical significance was confirmed by low Wilk's Lambda values (0.004 and 0.13, respectively, $p =$
440 0.000) were responsible for the clusterization of cuttlefish in the bidimensional score plot reported in Fig.
441 3A. Samples from French Atlantic were well discriminated from samples from Chioggia and Mediterranean
442 Sea by the DF1 and Chioggia samples well discriminated from Mediterranean samples by the DF2. The most
443 important loadings for the DF1 were Sr and Na, while Co was found to be the most influent element for the
444 DF2 (Fig. 3A). Based on these results, 100% of cuttlefish were correctly recognized in CV, but, when
445 performing the external test set validation, two samples from Mediterranean area were misclassified as
446 samples from French Atlantic and one French Atlantic sample as of Mediterranean origin, thus resulting in
447 an overall accuracy of the model of 97% (Table 3). Fisher's discriminant function coefficients for each
448 geographical provenance are reported in Supplementary Material, Table S5.

449 In S-LDA the forward selection method based on Wilk's Lambda values was chosen to select the
450 discriminating variables, taking 3.84 as the minimum partial F value to include a variable and 2.71 as the
451 maximum F value to exclude a variable from the model (see Supplementary materials, Supplementary
452 Experimental Section). Based on this, two DFs and 12 out of 52 discriminating elements were finally included
453 into the model, corresponding to Na, Co, B, K, Cd, V, U, Rb, Ni, Ba, Cu, and As (Supplementary Materials,
454 Table S6), of which Na, Co, B, Cd, U, and Cu were the same as those selected in LW-LDA.

455 The DF1 and DF2 enclosed 82.5% and 17.5% of variance of original data, with a canonical correlation of
456 97.4% and 88.5% respectively. The significant discriminatory ability of each DF was confirmed not only by

457 Wilk's Lambda values of 0.03 for the DF1 and 0.114 for the DF2 ($p = 0.000$), but also by 99% accuracy in
458 LOOCV (one Mediterranean sample erroneously classified as Atlantic sample) and 100% accuracy in
459 external validation (Table 3). Fisher's coefficients for each provenance are reported in Supplementary
460 Materials, Table S6.

461 Optimal classification results (100% sensitivity, specificity, and accuracy, Table 3) were obtained by the
462 application of PLS-DA when 21 elements were selected on the basis of their VIP scores ($VIP > 1$) and used
463 as classificatory variables (see Supplementary Materials, Table S7). Among these elements, ten were found
464 to be the same of those previously included in stepwise-LDA. Although Sr, Mn, Mo, Li, Ca, Mg, Se, Bi, Cs,
465 P, and Y were selected in addition, some of these (Sr, Mo, Li, Ca, and Mg) were shared with LW-LDA. The
466 highest VIP scores were 1.672 for V and 1.628 for Co, thus indicating the maximum contribution of these
467 elements to the geographical separation between cuttlefish.

468 The VIP-PLS-DA model was created by using six LVs explaining 90.1% of variance ($R^2X = 0.901$), of
469 which 92.2% was directly correlated with labels of groups ($R^2Y = 0.922$). In addition, the overall training
470 model's predictive power of 88.8% ($Q^2 = 0.888$), resulted in 100% of samples to be correctly classified both
471 in LOOCV and external validation (Table 3), without any possible misleading interpretation deriving from
472 overfit or overprediction of the model as assessed by permutation test (average R^2Y -y-intercept = 0.048;
473 average Q^2 -y-intercept = -0.371). Further information about this validation is reported in Supplementary
474 Materials, Supplementary Fig. S4.

475 As for the RMSECV and RMSEP values, these were found to be low (0.156 and 0.159, respectively) and
476 very close to each other, thus remarking the outstanding ability of the simplified model in categorizing
477 cuttlefish.

478 Fig. 3B shows the bidimensional score and loading graphs of the first two DFs obtained by S-LDA, while the
479 first two LVs obtained by VIP-PLS-DA are reported in Fig. 3C. In both cases, cuttlefish were well separated
480 from each other, with samples from Chioggia distributing in the lower right quadrant of the score plots. In
481 addition, both the DF1 and the LV1 were mainly responsible for sample discrimination in the score plot.

482 By looking at the loading plots, Chioggia cuttlefish were found to be positively correlated with U, V, Ni, Ba,
483 As, Rb, and K in S-LDA (Fig. 3B) and with Cs, Rb, K and P in PLS-DA (Fig. 3C). The highest loading
484 scores for Mediterranean cuttlefish were instead observed for Cd in S-LDA, which in turn was diagonally

485 opposite to the elements that characterized Chioggia samples. In PLS-DA model, the highest degree of
486 correlation with Mediterranean cuttlefish was instead found for V, Se, Co, Cu, Mo, Ni, Bi, and Mn while for
487 French Atlantic cuttlefish the highest contribution was exerted by Na together with Mg, U, Li, Sr, Ca and Cd.
488 Likewise, Na showed a high loading value on the DF2 negative axis and, therefore, it allowed to distinguish
489 French samples from the other groups in the S-LDA model.

490 According to the results of multi-element analysis of fish and seafood products already published in
491 literature, some of the most discriminant elements found in the present study might be linked to
492 anthropogenic pollution (As, V, Cd and U) or to geogenic sources (Cs and Mn) (Costas-Rodríguez et al.,
493 2010).

494 Consistently to the results achieved, the major elements Na, Mg, and K were also previously identified
495 among the most useful indicators of geographical origin of crabs (Luo et al., 2019) and prawns (Gopi et al.,
496 2019). The variation in the amounts of Na, Mg and P in the tissues of Pacific shrimps was also correlated
497 with specific sampling sites, whose waters were characterized by higher salinity (Li, Han, Dong, & Boyd,
498 2019).

499 The concentrations of As which, in the present work, was one of the selected discriminating elements for
500 Chioggia cuttlefish, were also reported in mussels from Venice Lagoon (in which Chioggia is located) as
501 being positively associated to the higher degree of salinity of this area and thus valuable for the identification
502 of local products (Cubadda, Raggi, & Coni, 2006). Moreover, also Ni, Co, Se, Mo, and V amounts showed a
503 strong link with anthropic sources of element contamination of the Venice Lagoon (Cubadda et al., 2006),
504 therefore the specific pattern distribution of these trace elements in Chioggia cuttlefish tissues might be
505 supposed to specifically reflect their origin.

506 Based on these results, the use of multivariate classifiers in combination with the pre-selection of the most
507 origin-discriminant elements, was proved to be an efficient, rapid and smart analytical strategy to assure the
508 authenticity of the provenance of cuttlefish samples. Although each of the classification techniques applied
509 showed satisfying results, suggesting that possible users may choose the most convenient methodology to
510 suit the specific needs, superior and unequivocal outcomes were obtained by applying VIP-PLS-DA.

511 **4. Conclusions**

512 Cephalopods are among some of the most consumed fishery products and are appreciated by the consumers
513 all over the world but, to the best of our knowledge, scientific insights concerning their elemental profile and
514 the authentication of their geographical origin are still lacking.

515 The results presented in this study revealed for the first time that the geographical imprint of Italian
516 traditional cuttlefish from Chioggia can be extracted through the simultaneous quantification of more than
517 fifty elements by ICP-MS and successfully used to discriminate the product from cuttlefish originating from
518 different areas.

519 The whole elemental profile was elaborated by means of different chemometric techniques. In particular,
520 three independent variable selection strategies merged with linear or regression pattern recognition
521 multivariate techniques (LW-LDA, S-LDA, and VIP-PLS-DA) were tested to develop classification rules
522 while reducing the number of variables to the lowest possible, in order to find the most parsimonious way
523 that best described sample origins. Although the application of VIP-PLS-DA led to the extraction of the
524 highest number of variables, analysis of performance metrics suggested the best results for this methodology,
525 since values of sensitivity, specificity, and classification accuracy of 100% were achieved in internal and in
526 external validation thanks to the contribution of Na, K, Ca, P, Mg, Cu, Co, Mn, Se, Ni, Mo, Li, B, Ba, Bi, Sr,
527 As, V, Rb, Cs, Y, U, and Cd.

528 In summary, the elemental pattern linked to the geographical origin of cuttlefish appears to be determined by
529 a combination of macro, trace and ultra-trace elements of both natural and anthropogenic origin, which are
530 known to be absorbed by the animals from the surrounding environment. In particular, the contribution of
531 anthropogenic elements here strongly emerges as a key analytical determinant for cephalopods authenticity
532 assessment. Of note, also concentrations of some heavy toxic metals such as Cd and As, although being so
533 low as not to represent a potential health risk, were useful for the discrimination purpose.

534 Considering the permanent and continuous release of all these elements in the aquatic environment deriving
535 from modern industrial and agricultural practices, the quantification of anthropogenic contaminants in
536 fishery products should therefore be encouraged, as well as further specific critical evaluations of their
537 variation within seawater is recommended. In such a scenario, the promising results obtained may pave the
538 way for practical application of the proposed methodology in the fishery sector to trace back different

539 geographical origins, but also for the protection and ongoing promotion of traditional local fishery products
540 for which a quality mark has been recognized.

541 **Appendix A. Supplementary Materials**

542 Supplementary data associated with this article can be found, in the online version, at

543 **Declaration of interest:** The authors declare that they have no known competing financial interests or
544 personal relationships that could have appeared to influence the work reported in this paper.

545 **Data Availability:** The dataset generated during the current study is available from the corresponding author
546 Prof. Lenka Husáková on reasonable request.

547 **Acknowledgements**

548 The authors gratefully acknowledge the financial support from the University of Pardubice (project no.
549 SGS_2020_002) and Dr. Giovanni Muresu Ibbi (DAVIMAR S.r.l, Chioggia, Italy) for providing cuttlefish
550 samples.

551 **References**

- 552 Abbas, O., Zadavec, M., Baeten, V., Mikuš, T., Lešić, T., Vulić, A., Prpić, J., Jemeršić, L., & Pleadin, J.
553 (2018). Analytical methods used for the authentication of food of animal origin. *Food Chemistry*, 246,
554 6–17. <https://doi.org/10.1016/j.foodchem.2017.11.007>
- 555 Arbuckle, N. S. M., & Wormuth, J. H. (2014). Trace elemental patterns in Humboldt squid statoliths from
556 three geographic regions. *Hydrobiologia*, 725(1), 115–123. <https://doi.org/10.1007/s10750-013-1608-4>
- 557 Ayas, D., & Ozogul, Y. (2011). The Effects of Season and Sex in the Metal Levels of Mature Common
558 Cuttlefish (*Sepia officinalis*) in Mersin Bay, Northeastern Mediterranean, *Journal of Food Science*,
559 76(4), 121–124. <https://doi.org/10.1111/j.1750-3841.2011.02152.x>
- 560 Bizzi, C. A., Pedrotti, M. F., Silva, J. S., Barin, J. S., Nóbrega, J. A., & Flores, E. M. M. (2017). Microwave-
561 assisted digestion methods: Towards greener approaches for plasma-based analytical techniques.
562 *Journal of Analytical Atomic Spectrometry*, 32(8), 1448–1466. <https://doi.org/10.1039/c7ja00108h>
- 563 Bosch, A. C., O'Neill, B., Sigge, G. O., Kerwath, S. E., & Hoffman, L. C. (2015). Heavy metals in marine
564 fish meat and consumer health: a review. *Journal of the Science of Food and Agriculture*, 96, 32–48.
565 <https://doi.org/10.1002/jsfa.7360>
- 566 Bua, G. D., Albergamo, A., Annuario, G., Zammuto, V., Costa, R., & Dugo, G. (2017). High-Throughput
567 ICP-MS and Chemometrics for Exploring the Major and Trace Element Profile of the Mediterranean
568 Sepia Ink. *Food Analytical Methods*, 10(5), 1181–1190. <https://doi.org/10.1007/s12161-016-0680-6>
- 569 Bustamante, P., Lahaye, V., Durnez, C., Churlaud, C., & Caurant, F. (2006). Total and organic Hg
570 concentrations in cephalopods from the North Eastern Atlantic waters: Influence of geographical origin
571 and feeding ecology. *Science of the Total Environment*, 368(2–3), 585–596.
572 <https://doi.org/10.1016/j.scitotenv.2006.01.038>
- 573 Carvalho, M. L., Santiago, S., & Nunes, M. L. (2005). Assessment of the essential element and heavy metal
574 content of edible fish muscle. *Analytical and Bioanalytical Chemistry*, 382, 426–432.
575 <https://doi.org/10.1007/s00216-004-3005-3>
- 576 Chaguri, M. P., Maulvault, A. L., Nunes, M. L., Santiago, D. A., Denadai, J. C., Fogaça, F. H., Sant'Ana, L.
577 S., Ducatti, C., Bandarra, N., Carvalho, M. L., Marques, A. (2015). Different tools to trace geographic

578 origin and seasonality of croaker (*Micropogonias furnieri*). *LWT - Food Science and Technology*,
579 61(1), 194–200. <https://doi.org/10.1016/j.lwt.2014.11.006>

580 Clodia database (2017). Database of Fishery Data from Chioggia, Northern Adriatic Sea.
581 <http://chioggia.biologia.unipd.it/banche-dati/>. Accessed 10.03.21

582 Commission of the European Communities (2006). Commission regulation (EC) No 1881/2006 of 19
583 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Official Journal of the*
584 *European Union*, L364/5 (20.12.2006). Retrieved from [https://eur-lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32006R1881)
585 [content/EN/ALL/?uri=CELEX%3A32006R1881](https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32006R1881)

586 Costas-Rodríguez, M., Lavilla, I., & Bendicho, C. (2010). Classification of cultivated mussels from Galicia
587 (Northwest Spain) with European Protected Designation of Origin using trace element fingerprint and
588 chemometric analysis. *Analytica Chimica Acta*, 664(2), 121–128.
589 <https://doi.org/10.1016/j.aca.2010.03.003>

590 Cubadda, F., Raggi, A., & Coni, E. (2006). Element fingerprinting of marine organisms by dynamic reaction
591 cell inductively coupled plasma mass spectrometry. *Analytical and Bioanalytical Chemistry*, 384(4),
592 887–896. <https://doi.org/10.1007/s00216-005-0256-6>

593 Danezis, G. P., Tsagkaris, A. S., Camin, F., Brusica, V., & Georgiou, C. A. (2016). Food authentication:
594 Techniques, trends & emerging approaches. *TrAC Trends in Analytical Chemistry*, 85, 123-132.
595 <https://doi.org/10.1016/j.trac.2016.02.026>

596 Drivelos, S. A., & Georgiou, C. A. (2012). Multi-element and multi-isotope-ratio analysis to determine the
597 geographical origin of foods in the European Union. *TrAC - Trends in Analytical Chemistry*, 40, 38–51.
598 <https://doi.org/10.1016/j.trac.2012.08.003>

599 European Food Safety Authority (EFSA), Panel on Contaminants in the Food Chain (CONTAM), (2009).
600 Scientific Opinion on Arsenic in Food. *EFSA Journal*, 7(10), 1351.
601 <https://doi.org/10.2903/j.efsa.2009.1351>.

602 European Parliament and Council of the European Union. (2013). Regulation (EU) No 1379/2013 of the the
603 European Parliament and of the Council of 11 December 2013 on the common organization of the
604 markets in fishery and aquaculture products, amending Council Regulations (EC) No 1184/2006 and

605 (EC) No 1224/2009 and repealin Council Regulation (EC) No 104/2000. *Official Journal of the*
606 *European Union*, L354(28.12.2013), 1–21. Retrieved from [https://eur-lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1588369230953&uri=CELEX:32013R1379)
607 [content/EN/TXT/?qid=1588369230953&uri=CELEX:32013R1379](https://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1588369230953&uri=CELEX:32013R1379)

608 FAO (Food and Agriculture Organization of the United Nations) (2020). FAO Species Fact Sheets, *Sepia*
609 *officinalis* (Linnaeus, 1758). <http://www.fao.org/fishery/species/2711/en>. Accessed 06.03.20

610 FAO/WHO (Food and Agriculture Organization of the United Nations/World Health Organization)
611 (2004). Evaluations of the Joint FAO/WHO Expert Committee on Food Additives (JECFA). From
612 twenty-sixth to seventy-fourth meetings. [https://apps.who.int/food-additives-contaminants-jecfa-](https://apps.who.int/food-additives-contaminants-jecfa-database/search.aspx)
613 [database/search.aspx](https://apps.who.int/food-additives-contaminants-jecfa-database/search.aspx). Accessed 03.05.20

614 Gonzalez, M. H., Souza, G. B., Oliveira, R. V., Forato, L. A., Nóbrega, J. A., & Nogueira, A. R. A. (2009).
615 Microwave-assisted digestion procedures for biological samples with diluted nitric acid: Identification
616 of reaction products. *Talanta*, 79(2), 396–401. <https://doi.org/10.1016/j.talanta.2009.04.001>

617 Gopi, K., Mazumder, D., Sammut, J., Saintilan, N., Crawford, J., & Gadd, P. (2019b). Combined use of
618 stable isotope analysis and elemental profiling to determine provenance of black tiger prawns (*Penaeus*
619 *monodon*). *Food Control*, 95, 242–248. <https://doi.org/10.1016/j.foodcont.2018.08.012>

620 Husáková, L., Urbanová, I., Šrámková, J., Černohorský, T., Krejčová, A., Bednaříková, M., Frýdová E.,
621 Nedělková, I., & Pilařová, L. (2011). Analytical capabilities of inductively coupled plasma orthogonal
622 acceleration time-of-flight mass spectrometry (ICP-oe-TOF-MS) for multi-element analysis of food and
623 beverages. *Food Chemistry*, 129(3), 1287–1296. <https://doi.org/10.1016/j.foodchem.2011.05.047>

624 Husáková, L., Urbanová, I., Šídová, T., Cahová, T., Faltys, T., & Šrámková, J. (2015). Evaluation of
625 ammonium fluoride for quantitative microwave-assisted extraction of silicon and boron from different
626 solid samples. *International Journal of Environmental Analytical Chemistry*, 95(10), 922–935.
627 <https://doi.org/10.1080/03067319.2015.1070409>

628 Iguchi, J., Isshiki, M., Takashima, Y., Yamashita, Y., & Yamashita, M. (2014). Identifying the origin of
629 *Corbicula* clams using trace element analysis. *Fisheries Science*, 80(5), 1089–1096.
630 <https://doi.org/10.1007/s12562-014-0775-1>

631 Kang, X., Zhao, Y., Shang, D., Zhai, Y., Ning, J., & Sheng, X. (2018). Elemental analysis of sea cucumber

632 from five major production sites in China: A chemometric approach. *Food Control*, 94(March), 361–
633 367. <https://doi.org/10.1016/j.foodcont.2018.07.019>

634 Lall, S.P. (2002). The minerals. J.E. Halver, R.P. Hardy (Eds.), *Fish Nutrition* (3rd ed.), Academic Press,
635 London, pp. 260–308

636 Li, L., Boyd, C. E., & Sun, Z. (2016). Authentication of fishery and aquaculture products by multi-element
637 and stable isotope analysis. *Food Chemistry*, 194, 1238–1244.
638 <https://doi.org/10.1016/j.foodchem.2015.08.123>

639 Li, L., Han, C., Dong, S., & Boyd, C. E. (2019). Use of elemental profiling and isotopic signatures to
640 differentiate Pacific white shrimp (*Litopenaeus vannamei*) from freshwater and seawater culture areas.
641 *Food Control*, 95(August 2018), 249–256. <https://doi.org/10.1016/j.foodcont.2018.08.015>

642 Li, Y., Chen, C., Li, B., Sun, J., Wang, J., Gao, Y., Zhao, Y., Chai, Z. (2006). Elimination efficiency of
643 different reagents for the memory effect of mercury using ICP-MS. *Journal of Analytical Atomic*
644 *Spectrometry*, 21(1), 94–96. <https://doi.org/10.1039/b511367a>

645 Luo, R., Jiang, T., Chen, X., Zheng, C., Liu, H., & Yang, J. (2019). Determination of geographic origin of
646 Chinese mitten crab (*Eriocheir sinensis*) using integrated stable isotope and multi-element analyses.
647 *Food Chemistry*, 274(March 2018), 1–7. <https://doi.org/10.1016/j.foodchem.2018.08.104>

648 Ministerial Decree 5 June 2014. XIV Revisione dell'Elenco Nazionale dei Prodotti Agroalimentari
649 Tradizionali. *G.U. Serie Generale n.141 del 20-06-2014 - Suppl. Ordinario n. 48*, Ministero delle
650 Politiche Agricole Alimentari e Forestali, MIPAAF: Rome, Italy, 2014. Retrieved from
651 <https://www.gazzettaufficiale.it/eli/id/2014/06/20/14A04557/sg>

652 Neff, J. M. (2002). Chapter 3 - Arsenic in the Ocean. *Bioaccumulation in Marine Organisms, Effect of*
653 *Contaminants from Oil Well Produced Water* (1st ed.). Elsevier Science, pp. 57–78.
654 <https://doi.org/10.1016/B978-008043716-3/50004-X>

655 Noack, C. W., Dzombak, D. A., & Karamalidis, A. K. (2014). Rare earth element distributions and trends in
656 natural waters with a focus on groundwater. *Environmental Science and Technology*, 48(8), 4317–
657 4326. <https://doi.org/10.1021/es4053895>

658 Ortea, I., & Gallardo, J. M. (2015). Investigation of production method, geographical origin and species

659 authentication in commercially relevant shrimps using stable isotope ratio and/or multi-element
660 analyses combined with chemometrics: An exploratory analysis. *Food Chemistry*, 170, 145–153.
661 <https://doi.org/10.1016/j.foodchem.2014.08.049>

662 Raimundo, J., Pereira, P., Vale, C., & Caetano, M. (2005). Fe , Zn , Cu and Cd concentrations in the
663 digestive gland and muscle tissues of *Octopus vulgaris* and *Sepia officinalis* from two coastal areas in
664 Portugal. *Ciencias Marinas*, 31(1B), 243–251. <https://doi.org/10.7773/cm.v31i12.91>

665 Rashid, N. A., Hussain, W. S. E. C., Ahmad, A. R., & Abdullah, F. N. (2019). Performance of classification
666 analysis: A comparative study between PLS-DA and integrating PCA+LDA. *Mathematics and*
667 *Statistics*, 7(4), 24–28. <https://doi.org/10.13189/ms.2019.070704>

668 Smith, G. (2018). Step away from stepwise. *Journal of Big Data*, 5(1). [https://doi.org/10.1186/s40537-018-](https://doi.org/10.1186/s40537-018-0143-6)
669 0143-6

670 Smith, R. G., & Watts, C. A. (2009). Determination of the country of origin of farm-raised shrimp (family
671 penaeide) using trace metal profiling and multivariate statistics. *Journal of Agricultural and Food*
672 *Chemistry*, 57(18), 8244–8249. <https://doi.org/10.1021/jf901658f>

673 Storelli, M. M., Giacomini-Stuffler, R., Storelli, A., & Marcotrigiano, G. O. (2007). Cadmium and
674 mercury in cephalopod molluscs : Estimated weekly intake. *Food Additives and Contaminants*, 23(1),
675 25–30. <https://doi.org/10.1080/02652030500242023>

676 Varrà, M. O., Ghidini, S., Zanardi, E., Badiani, A., & Ianieri, A. (2019). Authentication of European sea bass
677 according to production method and geographical origin by light stable isotope ratio and rare earth
678 elements analyses combined with chemometrics. *Italian Journal of Food Safety*, 8(1).
679 <https://doi.org/10.4081/ijfs.2019.7872>

680 **Figure Captions**

681 **Fig. 1.** Pearson's correlation heat-map of the investigated elements.

682 **Fig. 2.** Biplot resulting from PCA applied to all the elements (A) and selected elements (B). The sample
683 surrounded by the red dotted line in the biplot indicate an outlier according to Hotelling's T^2 test (95%
684 confidence interval). Blue triangles: samples from Chioggia (FAO fishing area 37.2.1); red diamonds:
685 samples from Mediterranean Sea (FAO fishing area 37.1/37.2); yellow circles: samples from French Atlantic
686 (FAO fishing area 27.7.e).

687 **Fig. 3.** Comparison of score (left) and loading graphs (right) from LW-LDA (A), S-LDA (B) and VIP-PLS-
688 DA (C) for cuttlefish samples from different origins. Blue triangles: samples from Chioggia (FAO fishing
689 area 37.2.1); red diamonds: samples from Mediterranean Sea (FAO fishing area 37.1/37.2); yellow circles:
690 samples from French Atlantic (FAO fishing area 27.7.e).

691

692

693

694

695

696

697

698

Table 1

Agilent 7900 ICP-MS operating conditions.

Parameter	Setting		
ICP			
Plasma mode	General purpose		
Rf power (27 MHz) (W)	1550		
Sampling depth (mm)	8		
Plasma gas flow (L min ⁻¹)	15		
Auxiliary gas flow (L min ⁻¹)	0.9		
Nebulizer gas flow (L min ⁻¹)	1		
Nebulizer pump (rps)	0.1		
Spray chamber temperature (°C)	2		
Mass spectrometer	No gas mode	He mode	HEHe mode ^a
Extract 1 (V)		0	
Extract 2 (V)		-250	
Omega bias (V)	-100	-120	-120
Omega lens (V)	9.7	7.8	9.6
Cell entrance	-30	-40	-140
Cell exit	-50	-60	-150
Deflect (V)	11.6	1	-77
Plate bias	-35	-60	-150
Helium flow (mL min ⁻¹)	0	5	10
OctP bias	-8	-18	-100
OctP RF		200	
Energy discrimination (V)		5	
Number of elements	39 ^b	12 ^c	5 ^d
Acquisition			
Points per peak	1		
Replicates	3		
Sweeps/replicate	100		
Total acquisition time (s)	75		

^a HEHe mode - high energy helium mode; Monitored isotopes (integration time): ^b ⁷Li, ¹¹B, ²⁴Mg, ⁶⁶Zn, ⁸⁵Rb, ⁸⁸Sr, ⁸⁹Y, ⁹⁰Zr, ⁹⁵Mo, ¹⁰¹Ru, ¹⁰³Rh, ¹⁰⁵Pd, ¹¹¹Cd, ¹¹⁸Sn, ¹²¹Sb, ¹³³Cs, ¹³⁸Ba, ¹³⁹La, ¹⁴⁰Ce, ¹⁴¹Pr, ¹⁴⁶Nd, ¹⁴⁷Sm, ¹⁵³Eu, ¹⁵⁷Gd, ¹⁵⁹Tb, ¹⁶³Dy, ¹⁶⁵Ho, ¹⁶⁶Er, ¹⁷²Yb, ¹⁷⁵Lu, ¹⁷⁸Hf, ¹⁸⁵Re, ¹⁹⁵Pt, ²⁰⁵Tl, ²⁰⁶⁺²⁰⁷⁺²⁰⁸Pb, ²⁰⁹Bi, ²³²Th, ²³⁸U (all 0.1 s); ^c ²³Na (0.3 s), ²⁷Al (0.1 s), ³⁹K, ⁴⁴Ca (both 0.3 s), ⁵¹V (1 s), ⁵²Cr, ⁵⁵Mn, ⁵⁶Fe, ⁵⁹Co, ⁶⁰Ni, ⁶³Cu, ¹⁰³Rh (all 0.3 s); ^d ³¹P (0.1 s), ⁷⁵As, ⁷⁸Se (both 1 s), ¹⁰³Rh (0.3 s).

Table 2 Multi-elemental composition of cuttlefish from different geographical origins. **Elements are sorted according to decreasing median concentrations of cuttlefish from Chioggia.**

Element	Chioggia (n = 17*3)				Mediterranean Sea (n = 25*3)				French Atlantic (n = 26*3)			
	Mean ± ME	Median	Min	Max	Mean ± ME	Median	Min	Max	Mean ± ME	Median	Min	Max
Na	11164 ± 867	11235 ^a	8659	14187	14334 ± 486	14290 ^b	12015	16338	24632 ± 1332	24730 ^c	17932	30749
K	13471 ± 1067	13768 ^a	9113	16488	10648 ± 402	10547 ^b	8898	12400	8353 ± 531	8252 ^c	6196	12039
P	10697 ± 831	10635 ^a	7459	1617	9084 ± 221	9079 ^b	7440	9883	8203 ± 308	8176 ^c	6791	10401
Mg	2297 ± 207	2274 ^a	1626	3180	2613 ± 75	2643 ^a	2164	2871	3767 ± 183	3776 ^b	2898	4795
Ca	820 ± 102	751 ^a	619	1338	1299 ± 53	1304 ^b	1019	1513	1781 ± 118	1760 ^c	1239	2493
As	79 ± 9	75.8 ^a	56.1	121.3	69 ± 4.3	67.4 ^{ab}	50.5	98.8	60 ± 4.7	62.3 ^b	35.5	77.1
Li	0.25 ± 0.024	0.24 ^a	0.18	0.34	0.34 ± 0.011	0.33 ^b	0.28	0.38	0.50 ± 0.025	0.51 ^c	0.36	0.65
Zn	62 ± 5.2	60.7 ^a	41.1	86.3	73 ± 1.6	72.8 ^b	63.8	80.7	74 ± 3.5	73.8 ^b	58.6	94.1
Al	16 ± 5.5	10.7 ^a	4.7	43.8	7 ± 1.9	5.69 ^b	2.97	22.7	7 ± 2.5	5.44 ^b	2.87	34.4
Cu	10 ± 1.8	8.55 ^a	5.93	20.2	24 ± 3.4	21.3 ^b	12.5	42.9	20 ± 3.4	16.3 ^b	8.29	36.2
Sr	9 ± 1.1	8.83 ^a	6.69	14.9	14.8 ± 0.44	14.9 ^b	12.0	16.8	25 ± 1.6	26.0 ^c	16.3	33.2
Fe	11 ± 4	7.85 ^a	3.62	33.0	9 ± 1.9	8.19 ^a	4.32	25.0	8 ± 2	5.59 ^a	3.41	21.7
B	6.3 ± 0.56	6.06 ^{ab}	4.46	8.20	5.3 ± 0.29	5.35 ^a	4.11	7.15	7.6 ± 0.85	7.88 ^b	4.15	11.8
Rb	5.3 ± 0.42	5.17 ^a	3.94	6.99	4.5 ± 0.14	4.61 ^a	3.93	5.13	3.6 ± 0.22	3.57 ^b	2.56	5.27
Se	1.3 ± 0.17	1.33 ^a	0.881	2.11	2.0 ± 0.14	1.91 ^b	1.47	2.69	1.65 ± 0.083	1.62 ^c	1.26	2.16
Mn	2.2 ± 0.87	1.28 ^a	0.93	6.86	2.5 ± 0.52	2.04 ^a	1.41	6.59	1.0 ± 0.23	0.83 ^b	0.54	3.57
Ni	0.4 ± 0.52	0.36 ^a	0.22	0.76	0.52 ± 0.056	0.50 ^b	0.27	1.98	0.28 ± 0.034	0.26 ^a	0.15	0.48
Cr	0.18 ± 0.060	0.18 ^a	0.037	0.55	0.15 ± 0.041	0.13 ^a	0.068	0.54	0.22 ± 0.082	0.13 ^a	0.081	1.07
Ba	0.3 ± 0.13	0.18 ^a	0.087	1.17	0.22 ± 0.031	0.16 ^a	0.12	0.50	0.21 ± 0.039	0.18 ^a	0.11	0.43
Pb	0.10 ± 0.024	0.093 ^a	0.050	0.23	0.10 ± 0.019	0.099 ^a	0.063	0.31	0.09 ± 0.017	0.046 ^a	0.077	0.20
V	0.07 ± 0.036	0.040 ^a	0.018	0.26	0.25 ± 0.040	0.22 ^b	0.12	0.47	0.11 ± 0.023	0.11 ^a	0.039	0.27
Zr	0.04 ± 0.018	0.027 ^a	0.0059	0.13	0.05 ± 0.015	0.036 ^{ab}	0.0071	0.14	0.2 ± 0.20	0.051 ^b	0.0066	2.55
Co	0.04 ± 0.019	0.025 ^a	0.016	0.15	0.21 ± 0.023	0.21 ^b	0.11	0.31	0.14 ± 0.026	0.14 ^c	0.058	0.31
Mo	0.029 ± 0.0066	0.024 ^a	0.018	0.060	0.10 ± 0.013	0.089 ^b	0.061	0.18	0.10 ± 0.019	0.084 ^b	0.044	0.23
Pd	0.016 ± 0.0013	0.016 ^a	0.012	0.020	0.024 ± 0.0054	0.021 ^a	0.012	0.067	0.029 ± 0.0033	0.027 ^b	0.017	0.059
Cs	0.017 ± 0.0015	0.016 ^a	0.012	0.023	0.0146 ± 0.00085	0.014 ^a	0.012	0.023	0.012 ± 0.0014	0.012 ^b	0.0081	0.027
Y	0.02 ± 0.012	0.015 ^a	0.0057	0.11	0.008 ± 0.0012	0.0074 ^b	0.0039	0.017	0.01 ± 0.013	0.0077 ^b	0.0050	0.17
Bi	0.178 ± 0.0022	0.017 ^a	0.011	0.027	0.023 ± 0.0040	0.021 ^a	0.015	0.065	0.010 ± 0.0041	0.0066 ^b	0.0029	0.044
Sn	0.015 ± 0.0071	0.011 ^a	0.0044	0.061	0.022 ± 0.0049	0.019 ^b	0.0075	0.051	0.015 ± 0.0040	0.011 ^a	0.0052	0.053
Cd	0.02 ± 0.017	0.0099 ^a	0.0035	0.13	0.08 ± 0.013	0.071 ^b	0.035	0.15	0.25 ± 0.059	0.22 ^c	0.067	0.58
Sb	0.007 ± 0.0028	0.0053 ^a	0.0023	0.021	0.009 ± 0.0012	0.0084 ^b	0.0046	0.016	0.0075 ± 0.00080	0.0068 ^b	0.0052	0.014
U	0.0032 ± 0.00076	0.0026 ^a	0.0019	0.0074	0.0043 ± 0.00042	0.0040 ^a	0.0027	0.0069	0.010 ± 0.0015	0.0091 ^b	0.0049	0.019
Th	0.002 ± 0.0011	0.0016 ^a	0.00081	0.0082	0.005 ± 0.0036	0.0022 ^a	0.0011	0.046	0.003 ± 0.0019	0.0015 ^a	0.0011	0.024
Ce*	29 ± 10	20.6 ^a	3.56	82.0	44 ± 20	24.0 ^a	11.4	250	26 ± 8.9	18.8 ^a	8.31	91.7
La*	20 ± 11	11.9 ^{ab}	2.70	90.87	29 ± 12	20.3 ^a	8.03	150	16 ± 5.6	10.5 ^b	5.09	56.0
Nd*	8 ± 4.7	3.98 ^a	1.58	33.2	5 ± 1.5	4.45 ^a	1.74	20.1	4 ± 1.0	2.82 ^b	1.33	12.0
Hf*	2.5 ± 0.52	2.23 ^a	1.15	4.94	8 ± 8.4	2.89 ^{ab}	1.42	103	8 ± 4.9	4.19 ^b	0.92	57.3
Tl*	1.5 ± 0.11	1.42 ^a	1.07	1.94	1.4 ± 0.51	1.21 ^b	0.83	7.30	1.3 ± 0.61	1.05 ^b	0.62	8.66
Pr*	2 ± 1.0	0.96 ^a	0.38	7.43	1.2 ± 0.34	0.97 ^a	0.45	4.62	0.8 ± 0.23	0.63 ^b	0.33	2.56
Sm*	1.5 ± 0.81	0.74 ^a	0.25	5.04	0.9 ± 0.28	0.82 ^a	0.34	3.73	0.7 ± 0.18	0.55 ^a	0.33	2.32
Dy*	1.1 ± 0.44	0.74 ^a	0.34	3.22	0.8 ± 0.18	0.66 ^{ab}	0.27	2.21	0.7 ± 0.17	0.51 ^b	0.33	2.02
Gd*	1 ± 0.5	0.69 ^{ab}	0.24	3.20	0.8 ± 0.19	0.72 ^a	0.32	2.52	0.6 ± 0.15	0.50 ^b	0.31	1.97
Er*	0.7 ± 0.23	0.54 ^a	0.20	1.73	0.5 ± 0.10	0.42 ^{ab}	0.23	1.17	0.4 ± 0.11	0.39 ^b	0.19	1.31
Yb*	0.7 ± 0.21	0.51 ^a	0.22	1.48	0.39 ± 0.078	0.37 ^{ab}	0.14	0.93	0.4 ± 0.10	0.34 ^b	0.16	1.31
Eu*	0.4 ± 0.18	0.24 ^{ab}	0.12	1.19	0.28 ± 0.061	0.25 ^a	0.15	0.87	0.22 ± 0.044	0.19 ^b	0.12	0.62
Ho*	0.22 ± 0.078	0.17 ^a	0.062	0.60	0.15 ± 0.030	0.14 ^{ab}	0.047	0.39	0.14 ± 0.037	0.11 ^b	0.063	0.45
Tb*	0.20 ± 0.082	0.14 ^a	0.056	0.58	0.15 ± 0.037	0.12 ^{ab}	0.055	0.43	0.12 ± 0.030	0.095 ^b	0.054	0.37
Ru*	0.10 ± 0.029	0.087 ^a	0.031	0.24	0.14 ± 0.027	0.13 ^a	0.054	0.34	0.14 ± 0.023	0.13 ^a	0.044	0.29
Re*	0.10 ± 0.032	0.081 ^{ab}	0.016	0.24	0.12 ± 0.019	0.11 ^a	0.070	0.24	0.09 ± 0.023	0.072 ^b	0.025	0.28
Lu*	0.09 ± 0.032	0.074 ^a	0.023	0.23	0.06 ± 0.015	0.047 ^a	0.021	0.18	0.06 ± 0.017	0.053 ^a	0.021	0.22
Pt*	0.13 ± 0.034	0.10 ^a	0.060	0.25	0.2 ± 0.12	0.12 ^a	0.062	1.53	0.15 ± 0.061	0.10 ^a	0.019	0.60
Hg [#]	0.20 ± 0.030	0.18 ^a	0.15	0.34	0.20 ± 0.023	0.17 ^a	0.14	0.38	0.21 ± 0.018	0.20 ^a	0.13	0.29

Concentrations are expressed as mg kg⁻¹ (d.m.) and are reported as mean values ± margin of error (ME) at 95% confidence level. Min–Max: minimum and maximum values found. *: concentrations are expressed as µg kg⁻¹. Hg[#]: determined by direct mercury analyzer AMA254. Values followed by different superscript letters (a–c) in the same row are significantly different ($p \leq 0.05$).

1 **Multi-element signature of cuttlefish and its potential for the discrimination of**
2 **different geographical provenances and traceability**

3

4 *Maria Olga VARRÀ^a, Lenka HUSÁKOVÁ^{b*}, Jan PATOČKA^b, Sergio GHIDINI^a,*
5 *Emanuela ZANARDI^{a*}*

6

7 *^aDepartment of Food and Drug, University of Parma, Parma, Via del Taglio, 10, Parma 43126, Italy*

8 E-mail addresses: mariaolga.varra@studenti.unipr.it (M. O. Varrà); sergio.ghidini@unipr.it (S.
9 Ghidini).

10 *^bDepartment of Analytical Chemistry, Faculty of Chemical Technology, University of Pardubice,*
11 *Studentska 573 HB/D, Pardubice, CZ-532 10, Czech Republic*

12 E-mail address: jan.patocka@upce.cz (J. Patočka)

13

14 ***CORRESPONDING AUTHORS:**

15 **Lenka Husáková** (L. Husáková)

16 Department of Analytical Chemistry, Faculty of Chemical Technology, University of Pardubice
17 Studentská 573 HB/D, Pardubice, CZ-532 10 (Czech Republic)

18 Tel. +420 466 037 029

19 Fax: +420 466 037 068

20 E-mail address: lenka.husakova@upce.cz

21

22 **Emanuela Zanardi** (E. Zanardi)

23 Department of Food and Drug, University of Parma, Parma, Strada del Taglio, 10, Parma 43126, Italy

24 Tel. +39.0521.902.760

25 E-mail address: emanuela.zanardi@unipr.it

26 **ABSTRACT**

27 The measurement and analysis of fifty-two elements by quadrupole inductively coupled plasma mass
28 spectrometry (Q-ICP-MS) and direct mercury analysis were applied to origin discrimination of Italian
29 traditional cuttlefish (Chioggia, Venice lagoon) from Mediterranean and Atlantic samples. A total 68
30 specimens were analyzed in triplicates to generate 204 mass spectra profiles which were statistically processed
31 by different chemometric techniques. Loading weights from principal component analysis as input for linear
32 discriminant analysis (LW-LDA), stepwise-LDA (S-LDA) and variable influence of projection-partial least
33 square discriminant analysis (VIP-PLS-DA) were used to classify samples while retaining the lowest possible
34 number of key variables. VIP-PLS-DA was found to be the best variable selection-discriminant tool combo
35 since the selected Na-Co-B-K-Cd-V-U-Rb-Ni-Ba-Cu-As-Sr-Mn-Mo-Li-Ca-Mg-Se-Bi-Cs-P-Y
36 elemental pattern allowed the samples to be classified with 100% sensitivity, specificity and accuracy.

37

38 **Abbreviations**

39 Certified reference materials, CRMs; correlation analysis, CA; discriminant function, DF; high energy Helium
40 mode, HE He; inductively coupled plasma mass spectrometry, ICP-MS; inductively coupled plasma optical
41 emission spectrometry ICP-OES; internal standard, ISTD; kinetic energy discrimination, KED; linear
42 discriminant analysis, LDA; leave-one-out-cross-validation, LOOCV; loading weights-based linear
43 discriminant analysis, LW-LDA; method detection limit of the method, MDL; method limit of quantification,
44 MLOQ; microwave digestion, MWD; partial least square-discriminant analysis, PLS-DA; principal component,
45 PC; principal component analysis, PCA; quadrupole inductively coupled plasma mass spectrometry, Q-ICP-
46 MS; rare earth elements, REEs; relative standard deviation, RSD; root mean square error from cross-validation,
47 RMSECV; root mean square error of estimation, RMSEE; root mean square error of prediction, RMSEP;
48 stepwise- linear discriminant analysis, S-LDA; variable influence on projection, VIP; variable influence of
49 projection-partial least square discriminant analysis, VIP-PLS-DA.

50

51

52 **Keywords:** ICP-MS; element profiling; fishery products; geographical origin; chemometrics.

53 **1. Introduction**

54 The common cuttlefish (*Sepia officinalis*, L., 1758) is a highly valuable fishery product whose distribution
55 extends from the Eastern Atlantic (from the Baltic and North Seas to South Africa coasts) up to the whole
56 Mediterranean Sea. Large scale fishing operations of common cuttlefish led to an average production of
57 21 825 000 kg in 2018, which ensured the product's supply all over the world (especially in Italy, Spain, Japan,
58 and South Korea), while smaller-scale fishing is particularly profitable in some Mediterranean areas where it
59 has a high impact on local economies (FAO, 2020). This activity is particularly developed in the fishing village
60 of Chioggia (north-western Adriatic Sea in the Eastern Mediterranean Basin), whose *S. officinalis* production
61 reached more than 454 000 kg in 2019 (Clodia database, 2017).

62 Chioggia cuttlefish is well known to be of high quality thanks to the preservation of a close link with the
63 territory and the traditional processing methods by local fish plants, which allow the products to be officially
64 certified as Italian Traditional Agri-food Product (P.A.T.) (Ministerial Decree, G.U. n. 48, 2014). Foodstuffs
65 with specific geographical ties (especially when a quality mark has been recognized) are in turn rated more
66 highly in terms of quality by the consumer, mostly because of lower environmental impact and higher
67 perceived safety of regional products.

68 Alongside with the compulsory labelling requirement established by European legislation concerning the
69 provision of detailed information about the geographical origin of fish and seafood (European Parliament and
70 Council of the European Union, 2013), the protection and the promotion of traditional foods must involve
71 origin authenticity assessment both to ensure traceability and prevent commercial frauds (Ortea & Gallardo,
72 2015).

73 Complex multi-disciplinary and cross-disciplinary approaches are usually required to verify the geographical
74 origin of fish and seafood since both environment and genetics can affect the final characteristics of the
75 products (Abbas et al., 2018). Nevertheless, considering the association between the concentrations of
76 elements in fish tissues and those in the surrounding aquatic environment, the determination of the multi-
77 element profile of fish and seafoods can be regarded as a valid analytical strategy to guarantee fish origin
78 authenticity. Several factors such as species, size, age, sex, and sexual maturity can directly affect the elemental
79 composition of aquatic animals, but food resources, climate, presence of contaminants, and many water quality

80 parameters (*pH*, dissolved oxygen, alkalinity) are those which differ to a greater extent between countries (Li,
81 Boyd, & Sun, 2016; Smith & Watts, 2009).

82 Several studies reported the possibility to discriminate fishery and seafood products as croaker (Chaguri et al.,
83 2015), sea cucumber (Kang et al., 2018), shrimps (Ortea & Gallardo, 2015), crabs (Luo et al., 2019), and
84 bivalve mollusks such as mussels (Costas-Rodríguez, Lavilla, & Bendicho, 2010) and clams (Iguchi, Isshiki,
85 Takashima, Yamashita, & Yamashita, 2014) using element composition. In these works, the chemical
86 characterization of samples was based on minor and/or trace elements (Costas-Rodríguez et al., 2010; Iguchi
87 et al., 2014), a combination of major, minor, and trace elements (Kang et al., 2018), or a combination of major
88 and trace elements plus stable isotopes analysis (Chaguri et al., 2015; Ortea et al., 2015; Luo et al., 2019).

89 In the range of analytical methods applied to the determination of elements, inductively coupled plasma-mass
90 spectrometry (ICP-MS) combined with different chemometrics techniques holds a unique position by virtue
91 of speed, sensitivity, dynamic range, and elemental coverage (Drivelos & Georgiou, 2012; Danezis, Tsagkaris,
92 Camin, Brusic, & Georgiou, 2016). In this context, several supervised and unsupervised chemometric tools
93 such as, principal component analysis (PCA) (Chaguri et al., 2015; Kang et al., 2018; Ortea et al., 2015),
94 cluster analysis and *k*-means hierarchical clustering (Ortea et al., 2015), linear discriminant analysis (LDA)
95 (Costas-Rodríguez et al., 2010; Iguchi et al., 2014; Ortea et al., 2015; Kang et al., 2018; Luo et al., 2019),
96 support-vector machine (Luo et al., 2019), soft-independent modelling of class analogy and artificial neural
97 networks (Costas-Rodríguez et al., 2010) were applied for authenticity and traceability purposes.

98 The determination of elemental composition of fish and seafood to trace their geographical provenance is
99 particularly suitable for cephalopods mollusks. Cephalopods such as cuttlefish are in fact at high trophic levels
100 in the aquatic food chain and the overall elements accumulated in tissues are good indicators of the surrounding
101 habitat (Bosch, O'Neill, Sigge, Kerwath, & Hoffman, 2015). In addition, cephalopods are non-migratory
102 animals and, therefore, their elemental composition can reasonably be expected to be constant during life (Gopi
103 et al., 2019). Nevertheless, very few studies addressed the topic of origin discrimination of cephalopods
104 mollusks using elemental composition, in spite of being focused on the analysis of a limited number of major
105 and trace elements included in hard structures such as statoliths (Arbuckle & Wormuth, 2014) or ink (Bua et
106 al., 2017). These components are in fact not always retained in the final commercial product and, therefore,
107 not always exploitable in practical food surveillance operations to trace back the origin of cephalopods.

108 Based on this background, the present work aimed at outlining for the first time the *S. officinalis* multi-
109 elemental profile of edible tissues to verify whether useful information could be extracted and specifically
110 linked to the geographical origin of the Italian traditional cuttlefish from Chioggia (FAO subarea 37.2.1) in
111 order to be differentiated from non-Chioggia cuttlefish caught in other fishing areas of the Mediterranean Sea
112 (FAO 37.1/37.2) and in the French Atlantic Ocean (FAO 27.7.e). For this purpose, fifty-one elements were
113 determined by ICP-MS and Hg was quantified via atomic absorption spectroscopy. Unsupervised and
114 supervised pattern recognition methods were used to investigate sample characteristics, identify the key
115 discriminant elements, and, at the same time, develop classification rules.

116 **2. Materials and Methods**

117 *2.1. Reagents and standards*

118 The ultrapure water ($0.055 \mu\text{S cm}^{-1}$ conductivity) obtained using the Milli-Q[®] water purification system
119 (Millipore, Bedford, USA) was used for the preparation of all solutions. Sub-boiled nitric acid was prepared
120 from nitric acid (65%, w/w) of Selectipur quality (Lach-Ner, Neratovice, Czech Republic) using the distillation
121 equipment BSB-939-IR (Berghof, Eningen, Germany). Hydrogen peroxide (Trace Select, $\geq 30\%$, w/w) was
122 purchased from Fluka Chemie AG (Buchs, Switzerland).

123 Multi-element stock solution “A” containing 10 mg L^{-1} of Li, B, Al, V, Cr, Fe, Ni, Co, As, Se, Rb, Sr, Zr, Mo,
124 Ru, Pd, Cd, Sn, Sb, Cs, Ba, Hf, Re, Pt, Tl, Pb, Bi, and Th was prepared from the Supelco ICP multi-element
125 standard solution IV (Merck, Darmstadt, Germany) and single element standards of concentration
126 $1 \pm 0.002 \text{ g L}^{-1}$ (Analytika Ltd., Prague, Czech Republic or SCP Science, Montreal, Canada). Multi-element
127 solution “B” containing 1 mg L^{-1} of La, Ce, Pr, Nd, U (“B1”) and 0.20 mg L^{-1} of Y, Tb, Ho, Yb, Sm, Eu, Gd,
128 Er, Lu, and Dy (“B2”) was prepared from the stock solution of rare earth elements Astatol mix “M008”
129 (Analytika Ltd., Prague, Czech Republic). Multi-element solution “C” containing 50 mg L^{-1} of Na, Mg, P, K,
130 Ca, Mn, Cu, and Zn was prepared from single element standards of 1 g L^{-1} obtained from Analytika Ltd. The
131 internal standard solution (ISTD) was prepared from 1 g L^{-1} stock solution of Rh obtained from SCP Science
132 (Montreal, Canada). Carbon reference solutions were prepared from 10 g L^{-1} of C stock solution prepared from
133 urea of TraceSelect quality (Fluka Chemie AG, Buchs, Switzerland).

134 *2.2. Sampling and preparation of cuttlefish*

135 A total of 68 samples of common cuttlefish (*S. officinalis*, L.) including n = 17 samples from Adriatic Sea
136 (Chioggia, Italy, FAO 37.2.1), n = 25 non-Chioggia samples from the Mediterranean Basin (FAO 37.1/37.2),
137 and n = 26 samples from North-eastern Atlantic Ocean (France, FAO 27.7.e) were analyzed. Specimens were
138 caught by fishing trawlers during the months of September and randomly collected from different batches in
139 local fish plants located in Chioggia (Venice, Italy), by choosing homogeneous sizes and weights (mantle
140 lengths from 10 ± 2 cm; body weight 125 ± 25 g). After collection, samples were immediately frozen and
141 stored at -20 °C for around 3 months prior to be further processed for multi-element analysis.

142 *2.3. Quality assurance and quality control*

143 The following commercially supplied certified reference materials (CRMs) were analyzed: NIST SRM 1577
144 Bovine Liver (National Institute of Science and Technology, NIST, Gaithersburg, MD, USA); NIST SRM
145 1566 Oyster Tissue (NIST, Gaithersburg, MD, USA); BCR[®] certified reference material (CRM)184 Bovine
146 muscle (Institute for Reference Materials and Measurements, IRMM, Geel, Belgium); BCR[®] 185 Bovine Liver
147 (IRMM, Geel, Belgium); CRM NCS ZC73015 Milk Powder (National Research Centre for Certified
148 Reference Materials, NRCRM, Beijing, China); P-WBF CRM 12-2-04 Essential and Toxic Elements in Wheat
149 Bread Flour (pb-anal, Kosice, Slovakia); CRM12-2-03 P-Alfalfa Essential and toxic elements in Lucerne (pb-
150 anal, Kosice, Slovakia); SMU CRM 12-02-01 Bovine liver (pb-anal, Kosice, Slovakia).

151 *2.4. Sample preparation*

152 *2.4.1. Pre-processing of cuttlefish*

153 Frozen samples were thawed overnight (16–18 h at $+4$ °C) before processing for multi-element analysis. Each
154 specimen was then washed with deionized water and skin, cuttlebone, gills, reproductive and digestive tracts
155 were carefully removed without causing their rupture. The head, arms and tentacles were excluded, while the
156 mantle and the lateral fins were rinsed again with ultrapure water and minced with a ceramic knife.

157 *2.4.2. Freeze-drying process*

158 Pre-treated sample portions of approx. 10 g were individually transferred into 100 mL cleaned round bottom
159 flasks wherein the material was dried and placed into a deep freezer at $-80\text{ }^{\circ}\text{C}$ for 24 hours to provide a
160 necessary conditioning for drying. Lyophilization was conducted at $-50\text{ }^{\circ}\text{C}$ and 0.01 bar for 24 h with a LIO-
161 5P apparatus (CinquePascal srl, Trezzano sul Naviglio, Milan, Italy). Dried samples were subsequently
162 removed from the flasks, homogeneously ground into a powder by using ceramic mortars and pestles and then
163 individually sealed into LDPE bags.

164 *2.4.3. Microwave assisted digestion*

165 Microwave digestion (MWD) of samples was carried out by using the SpeedwaveTM MWS-3+ (Berghof,
166 Eningen, Germany) microwave system with the maximum total output of the microwave generator (1450 W)
167 and equipped with the optical sensor technology for contactless real-time recording of the sample temperature.
168 The high-pressure resistant (up to 100 bar) PTFE vessels DAC-100S, together with the Multitube System (all
169 from Berghof, Eningen, Germany), were used for sample digestion. This arrangement allowed the
170 simultaneous digestion of three samples in one DAC-100S PTFE vessel by placing three PFA tubes into each
171 vessel (Husáková et al., 2015). The maximal number of used DAC-100S vessels for one digestion cycle was
172 eight.

173 A powdered cuttlefish or CRM sample aliquot of 0.1 g was weighed into a 10 mL PFA tube. Then, 4 mL of
174 16% HNO_3 (65%, w/w HNO_3 , 1:3 diluted) and 1 mL of 30% H_2O_2 were added, leaving the vessels open until
175 the initial reaction subsided. Three PFA tubes containing the same sample and reagents were placed into the
176 outer 100mL PTFE digestion vessel previously filled with 25 mL of HNO_3 (16%, v/v), by ensuring that the
177 level of liquid in the outer PTFE vessel was higher than those in the PFA tubes. This way, the vapor pressures
178 were compensated and the evaporation of the solution from the PFA tubes was avoided (Husáková et al., 2015).
179 The samples were digested following a five-step program: (i) 20 min at $180\text{ }^{\circ}\text{C}$ and 80 % power (ramp 5 min),
180 (ii) 20 min at $220\text{ }^{\circ}\text{C}$ and 95 % power (ramp 5 min), (iii–v) 5 min at $100\text{ }^{\circ}\text{C}$ and 10 % power (ramp 1 min).
181 The resulting colorless solutions were diluted to 25 mL with deionized water.

182 Each sample was mineralized in three replicates. Blanks, consisting of deionized water and reagents were
183 subjected to a similar preparation procedure.

184 To assess the properness of the digestion process, the residual carbon content in the final cuttlefish digests was
185 determined by a previously described inductively coupled plasma optical emission spectrometry (ICP-OES)
186 method (Husáková et al., 2011) was $5.5 \pm 0.4 \%$ ($n = 3$).

187 2.5. ICP multi elemental analysis

188 ICP-MS measurements were performed by using the Agilent 7900 quadrupole mass spectrometer (Q-ICP-MS,
189 Agilent Technologies, Inc., Santa Clara, CA, USA) equipped with an octopole-based collision cell for
190 interference removal using kinetic energy discrimination (KED).

191 The standard sample introduction system, consisting of a glass concentric nebulizer MicroMist ($400 \mu\text{L min}^{-1}$)
192 ¹), the Peltier-cooled ($2 \text{ }^\circ\text{C}$) Scott quartz spray chamber and quartz torch with 2.5 mm internal diameter
193 injector, was used. For precise delivery of samples and ISTD, a low-pulsation, 10-roller peristaltic pump with
194 three separate channels was employed. The internal standard kit, including connecting tubing, connectors and
195 the “Y” piece was used for simultaneous internal standard aspiration and its mixing with the sample. The
196 standard sampling and skimmer nickel cones with orifices of 1 and 0.45 mm, respectively, were used. ICP-MS
197 operating conditions were optimized during each start-up sequence by using the multi-elemental tuning
198 solution (Agilent Technologies, Inc., Santa Clara, CA, USA) containing $1 \mu\text{g L}^{-1}$ of Ce, Co, Li, Mg, Tl and Y,
199 in order to obtain the highest possible sensitivity for elements of low, middle and high m/z . Using the typical
200 operating conditions summarized in Table 1, a sensitivity of $6000 \text{ counts s}^{-1} \text{ per } \mu\text{g L}^{-1}$ and a resolution of 0.64
201 amu peak width (full width at half maximum intensity) were achieved for $^7\text{Li}^+$. The same parameters were
202 $50000 \text{ counts s}^{-1} \text{ per } \mu\text{g L}^{-1}$ and 0.62 for $^{89}\text{Y}^+$, and $30000 \text{ counts s}^{-1} \text{ per } \mu\text{g L}^{-1}$ and 0.60 for $^{205}\text{Tl}^+$.

203 For sample analysis the “General Purpose” plasma mode included in the ICP-MS MassHunter software was
204 used (Agilent Technologies, Inc., Santa Clara, CA, USA). The working parameters of the cell mode “no-gas”
205 were autotuned during the instrument start-up sequence. The working parameters of the collision cell for
206 helium (“He”) and high energy He (“HE He”) modes were adjusted manually. The time required for a transition
207 between cell modes was 5 s. Parameters related to sample introduction and plasma conditions were consistent
208 for all modes (see Table 1).

209 Concentrations of a total of 51 elements were evaluated from calibration curves with coefficients of
210 determination better than 0.999 and built up within the ranges given below.

211 Calibration solutions: blank, 1, 5, 10, 50, 100 $\mu\text{g L}^{-1}$ of Li, B, Al, V, Cr, Fe, Ni, Co, As, Se, Rb, Sr, Zr, Mo,
212 Ru, Pd, Cd, Sn, Sb, Cs, Ba, Hf, Re, Pt, Tl, Pb, Bi, Th; 0.1, 0.5, 1, 5, 10 $\mu\text{g L}^{-1}$ La, Ce, Pr, Nd, U; 0.02, 0.1, 0.2,
213 1, 2 $\mu\text{g L}^{-1}$ of Y, Tb, Ho, Yb, Sm, Eu, Gd, Er, Lu, and Dy; 0.5, 1, 5, 10 mg L^{-1} of Na, Mg, P, K, Ca, Mn, Cu,
214 and Zn. The solutions were prepared daily by appropriate dilution of multi-element solutions “A” (500 $\mu\text{g L}^{-1}$),
215 “B” (50 + 10 $\mu\text{g L}^{-1}$) and “C” (50 mg L^{-1}) in 25 mL volumetric flasks (see Section 2.1).
216 To compensate possible instrumental drift and matrix effects, a 200 $\mu\text{g L}^{-1}$ Rh ISTD was simultaneously
217 aspired and mixed with samples.

218 2.6. Mercury analysis

219 For the quantification of Hg content, the direct solid sampling analysis using a single-purpose atomic
220 absorption spectrometer AMA 254 (Altec Ltd., Prague, Czech Republic), based on in situ dry ashing followed
221 by gold amalgamation atomic absorption spectroscopy, was used. Each time, samples were weighed in a nickel
222 boat and analyzed under the following experimental conditions: typical sample mass, 50 mg; drying time, 60 s;
223 decomposition, $\sim 750\text{ }^{\circ}\text{C}$ for 150 s; waiting step, $900\text{ }^{\circ}\text{C}$, 45 s, necessary for quantitative release of trapped
224 mercury from the gold amalgamator to the measuring cuvette. The peak area absorbance at 253.6 nm was
225 monitored. The flow rate of oxygen (99.5%) carrier gas was 170 mL min^{-1} .

226 The AMA 254 spectrometer was regularly calibrated using blank and four working standard solutions
227 containing Hg(II) ranging from 0.025 to 0.2 mg L^{-1} . The working standard solutions were prepared from
228 standard aliquots of 10 mg L^{-1} prepared from a 1 g L^{-1} stock solution of Hg(II) (Analytika Ltd., Prague, Czech
229 Republic).

230 2.7. Data treatment

231 Statistic was applied to elemental concentrations referring to cuttlefish dry matter (d.m.) and carried out using
232 IBM-SPSS (v. 23.0, SPSS Inc., Chicago, IL, USA) SIMCA-P v.14.1 (Umetrics, Umeå, Sweden), and
233 Statgraphics Centurion v.18.1 (StatPoint Technologies, Inc, Warrenton, VA, USA) software. The normality
234 and homogeneity of variance of data was checked beforehand by applying the Shapiro-Wilk's and Levene's
235 tests, respectively ($p \leq 0.05$). Since the normal distribution assumption was violated for most of the elemental
236 data, quantitative differences between groups of cuttlefish from different origins were investigated by the

237 nonparametric Kruskal–Wallis test with Dunn’s multiple post hoc test for multiple comparison ($p \leq 0.05$). Data
238 was presented as mean values \pm margin of error (calculated at 95% confidence level), median and minimum
239 and maximum values found.

240 While the nonparametric statistic was used to compensate for the lack of data normal distribution, thus
241 preserving the original information and avoiding the loose of any intuitive meanings when describing element
242 concentrations resulting from univariate data analysis, a Box-Cox transformation was instead applied to
243 normalize data for subsequent bivariate and multivariate data analyses.

244 First, the bivariate Pearson’s correlation analysis (CA) was employed to Box-Cox transformed data to explore
245 variable distribution and look for strong positive ($r > 0.6, p \leq 0.01$) or strong negative ($r < -0.6, p \leq 0.01$)
246 linear correlations between binary variables.

247 Because of the highly different magnitude of values among element concentrations, the Box-Cox data were
248 then scaled applying a Z-score standardization and further processed by multivariate data analyses. PCA was
249 used for further investigation of covariance patterns in elemental concentrations, data’s first interpretation and
250 multivariate outliers’ removal based on Hotelling’s T^2 test results (95% confidence interval). During this step,
251 the number of variables was also reduced by selecting elements characterized by loading values (rescaled to
252 $0-1) > 0.70$.

253 Afterwards, supervised classifiers combined with different variable selection procedures were developed to
254 achieve the discrimination of samples according to the geographical origin by using the lowest number and the
255 most influential combination of variables. To that end, linear discrimination based on variable loading weights
256 previously selected by PCA (LW-LDA), forward stepwise-LDA (S-LDA) and variable influence on
257 projection-(VIP)-PLS-DA were independently tested and compared in terms of prediction ability. All the
258 models were cross-validated using leave-one-out-cross-validation (LOOCV) to exclude overfitting. In addition
259 to LOOCV, the validity of the PLS discriminant models was also checked by permutation testing (400 random
260 permutations).

261 When supervised modelling was performed, about 33% of data ($n = 68$) was randomly sidelined from the
262 calibration set ($n = 136$) and used for independent external validation. Models’ overall performances in internal
263 and external validation were evaluated in terms of accuracy (%), sensitivity (%), and specificity (%).

264 Further details about the statistical treatments applied can be found in Supplementary material, Supplementary
265 Experimental Section.

266 **3. Results**

267 *3.1. Analytical performances and validation*

268 The analytical determination of fifty-two major, trace and ultra-trace elements in mineralized cephalopod
269 samples was carried out by Q-ICP-MS, taking advantage of its well-known sensitivity toward the targeted
270 elements. On the other hand, analysis of Hg by thermal decomposition was performed by AMA254 without
271 pre-treatment and/or preconcentration steps, thus overcoming the most serious problems related to ICP-MS
272 analysis of Hg, e.g. long washout time, non-linear calibration curves, and decreasing sensitivity with time (Li
273 et al., 2006).

274 The microwave-assisted pressurized digestion which involved small sample amounts (100 mg) was employed
275 to ensure rapid sample preparation for subsequent ICP-MS analysis while attaining the high decomposition
276 efficiency. One of the most important aspects regarding the employed MWD method is the possibility to use
277 diluted solutions for sample preparation, thus reducing the quantity of reagents, risks of contamination and
278 generation of residues. These are important parameters for the development of greener analytical methods
279 (Gonzalez et al., 2009). Moreover, MWD plays an important role in decreasing some kinds of spectral and
280 non-spectral interferences, such as those deriving from carbon containing species or different chlorides, by
281 conversion to significantly less or non-interfering species (Husáková et al., 2011; Bizzi et al., 2017). He mode
282 with KED or HE He mode with higher cell gas pressure and higher energy discrimination voltages, were used
283 for the analytes that suffered from spectral effects (i.e. V, Cr, Mn, Co, Ni, Fe, Co, Cu, As, and Se) caused by
284 polyatomic ions deriving from the plasma gas, sample solvent and other sample matrix elements such as Na,
285 K, Ca, Mg, P, S, C, and Cl (see Table 1 and Supplementary Materials, Table S1). He and/or HE He cell modes
286 had the additional benefit of reducing the response for low mass matrix elements like Na, K, Ca or P by an
287 order of magnitude ($^{23}\text{Na}^+$, $^{44}\text{Ca}^+$) or more ($^{39}\text{K}^+$, $^{31}\text{P}^+$), thus effectively raising the upper linear range for these
288 elements. Thereby elements that would have needed to be analyzed by ICP-OES, were included in the ICP-
289 MS.

290 As for the ISTD, Rh was chosen for its mid-range mass and ionization potential and because of its absence in
291 the analyzed samples. Recoveries of Rh, relative to the initial calibration blank for the entire 8-hour sequence
292 of sample digests, are shown in Supplementary Materials, Fig. S1. ISTD behavior across the mass range was
293 found to be very consistent over time and downward drift was considered acceptable.

294 Data accuracy was evaluated by the analysis of different CRMs (NIST SRM 1577 Bovine Liver, NIST 1566
295 Oyster Tissue, BCR CRM 184 Bovine muscle, BCR CRM 185 Bovine Liver, CRM 12-2-01 Bovine Liver),
296 primarily intended for the evaluation of analytical methods and instruments used for the determination of the
297 mass fraction values of selected elements in marine tissue, foods, or similar materials. Since the levels of most
298 lanthanides and actinides are not certified in these CRMs, three additional certified standards (NCS ZC 73015
299 Milk Powder, CRM 12-2-04 Wheat bread flour, and CRM 12-2-03 Essential and toxic elements in Lucerne)
300 were analyzed to validate data for these elements. The high level of agreement between target and found values
301 demonstrated trueness of data obtained (Supplementary Materials, Table S2 and Fig. S3).

302 The precision of the method was evaluated in terms of intra-day and inter-day comparison. Intra-day precision
303 was determined by analysis of individual control materials three times during the same day. Inter-day precision
304 was determined by analysis of the same standards on three different days over a period of one month. Within
305 each series, every solution was analyzed in triplicate. Relative standard deviation (RSD) was calculated for
306 both series of analyses. The RSD values of intra-day and inter-day studies were mostly found to be below 14%
307 thus showing a good precision of the method (Supplementary Materials, Table S2).

308 Method detection limits (MDLs) and method quantification limits (MLOQs) - reported in Supplementary
309 Material, Table S3 - were evaluated as a triple and tenfold standard deviation, respectively, of ten consecutive
310 measurements of blank signal divided by the slope of the calibration curve. Detection limits were found to be
311 low enough that selected elements could be determined at the background level. Fig. S2 and Table S3
312 (Supplementary Materials) also summarize relative sensitivities of Q-ICP-MS for analysis of individual
313 elements with the use of Rh ISTD.

314 *3.2. Concentrations of elements in cuttlefish samples*

315 The concentrations of elements measured in the present study were presented and discussed as median values
316 considering the non-normal distribution of the raw data. Results from descriptive statistics are listed in Table
317 2. The most abundant elements were found to be Na, Mg, P and K, whose concentration levels were higher

318 than 1% of weight and quite variable among cuttlefish from different countries. According to results from
319 Kruskal–Wallis test, Na and Mg amounts were significantly higher ($p \leq 0.05$) in samples from FAO 27.7.e
320 (French Atlantic) compared to samples from FAO 37.1/37.2 (Mediterranean area) and FAO 37.2.1 (Chioggia).
321 At the same time, an opposite trend was observed for P and K, for which the highest concentrations ($p \leq 0.05$)
322 were found in Chioggia cuttlefish (Table 2).

323 Concentrations between 1 and 0.01% of weight (10000–100 mg kg⁻¹, d.m.) were found for two elements
324 corresponding to Li and Ca, which showed significantly higher median concentrations in French Atlantic
325 samples (0.51 mg kg⁻¹ and 1760 mg kg⁻¹, respectively) ($p \leq 0.05$) compared to the other groups. The wide
326 variability in both major and minor element contents found in cuttlefish specimens of different geographical
327 origins and within samples of the same origin can be attributed to their natural variation in seawater, but the
328 assimilation by marine animals of Na, Mg, and Ca from the organic matter in the aquatic environment also
329 varies with the feeding status and the age of the animal (Lall, 2002). Moreover, the presence of P in fish and
330 seafood tissues can be directly attributable to the food sources since its concentration in seawater is lower
331 compared to the other elements (Lall, 2002).

332 The main trace elements, ranging between 100 and 1 mg kg⁻¹, followed the decreasing order Zn > Al > Sr >
333 Cu > Fe > B > Rb > Se > Mn in samples from Chioggia, Zn > Cu > Sr > Fe > Al > B > Rb > Mn > Se in
334 samples from the Mediterranean area, and Zn > Sr > Cu > B > Fe > Al > Rb > Se > Mn in samples from the
335 French Atlantic. Among these elements, only Se, Rb, and Sr were different in median concentrations according
336 to all three geographical regions ($p \leq 0.05$) and the concentration ranges were in line with those previously
337 published in literature for *S. Officinalis* (Raimundo, Pereira, Vale, & Caetano, 2005; Ayas & Ozogul, 2011).

338 Different median contents varying between 62.3 and 75.8 mg kg⁻¹ were also found for As ($p \leq 0.05$). The wide
339 presence of arsenic in the aquatic environment is linked both to the geochemical characteristics of the region
340 and anthropogenic activities, especially those related to agricultural practices (Neff, 2002). Due to toxic
341 properties, some concerns regarding the dietary exposure to arsenic have arisen in recent times, but maximum
342 levels for this metal in fishery products have not been established yet by European legislations. Fish and
343 seafood, in particular, have been reported as the largest food contributors to overall total arsenic exposure,
344 despite the largest proportion is represented by organic arsenic species which are known to be less or no toxic
345 compared to relative inorganic species (European Food Safety Authority, EFSA, 2009). In addition, inorganic

346 arsenic concentrations decrease with increasing content of total arsenic (European Food Safety Authority,
347 EFSA, 2009).

348 Most of the elements analyzed in the present paper were found to be present in cuttlefish tissues at mean
349 concentration lower than 1 mg kg⁻¹. Levels of ultra-trace elements as rare earth elements (REEs) were lower
350 than 100 µg kg⁻¹ on average, with the highest median concentration for Ce (24.0 µg kg⁻¹) and the lowest one
351 for Lu (0.047 µg kg⁻¹) both in in Mediterranean samples (Table 2). The natural pattern of REEs in the aquatic
352 environment, which is associated to the mobilization of nutrients from the underlying soils, can be permanently
353 altered by some anthropic activities and effectively used to investigate the geographical provenance of marine
354 animals (Noack, Dzombak, & Karamalidis, 2014). In the present work, the significantly different
355 concentrations of Tb, Dy, Ho, Er, and Yb geographically discriminated Chioggia from French specimens,
356 while those of La, Pr, Nd, Eu, and Gd discriminated Mediterranean from French specimens ($p \leq 0.05$). In
357 accordance with these results, the amount of both La and Ho was previously reported to be potentially useful
358 to authenticate fish samples from different areas in the Mediterranean Sea (Varrà, Ghidini, Zanardi, Badiani,
359 & Ianieri, 2019).

360 Regardless the origin, all the cuttlefish samples analyzed in the present work were in line with the maximum
361 limits for toxic heavy metals established by European regulations No. 1881/2006/EC and subsequent
362 amendments and additions (Commission of the European Communities, 2006). Both cadmium and lead did
363 not exceed the threshold value of 1 mg kg⁻¹ set for the edible part of cephalopods, as well as Hg was always
364 found to be below the threshold limit of 0.5 mg kg⁻¹. The sample groups under investigation did not show
365 significant differences in Pb and Hg concentrations ($p > 0.05$), but Cd median amount in French Atlantic
366 specimens was found to be approximately 7 and 20 times higher than in Mediterranean and Chioggia samples,
367 respectively (see Table 2). Furthermore, the comparison with results reported by other authors for cuttlefish
368 from North eastern Atlantic and Adriatic Sea, showed that Hg, Pb, and Cd concentrations found in the present
369 work fell within the same ranges (Bustamante, Lahaye, Durnez, Churlaud, & Caurant, 2006; Storelli,
370 Giacomini-Stuffler, Storelli, & Marcotrigiano, 2007). Even in the case of heavy toxic metals, a close link
371 between their concentrations in seawater and environmental pollution of specific areas exists (Carvalho,
372 Santiago, & Nunes, 2005). This, together with Ni, Co, and Bi amounts, which also varied significantly among

373 sample groups, makes suggestions for future insights about the effective role and contribution of these elements
374 for the geographical origin assessment of fish and seafood.

375 *3.3. Chemometric classification of the geographical origin of cuttlefish*

376 Since no specific information concerning the best arrangement of elements to discriminate the geographical
377 origin of cephalopods is available in literature, the first stages of multivariate analysis took into consideration
378 all the fifty-two elements measured, starting with the data exploration by means of CA and PCA. Nevertheless,
379 multicollinearity between variables as well as geographical-unrelated variability are a source of increasing
380 noise, thus the reduction of the variables to the most predictive ones was subsequently preferred, especially in
381 view of a future practical implementation of the methodology.

382 The selection of variables is strictly dependent on the sample matrix properties and dataset overall
383 characteristics; hence its application needs to be evaluated on an individual basis. Although being the most
384 frequently used classification technique in studies dealing with element profiling, S-LDA is
385 coming under growing criticism because of the deceiving results deriving from the random rather than
386 effective significance of the variables extracted, especially when the number of potential explanatory variables
387 is high (Smith, 2018). On the other hand, one of the main problems related to the use of multivariate data
388 analysis in food authentication studies is just the difficulty in comparing the obtained results with those already
389 published but using a wide variety of different chemometrics techniques. Therefore, in the context of the
390 present work, S-LDA and LW-LDA, were used for classification by variable selection, but outputs were
391 compared with those obtained through the application of VIP-based PLS-DA. The latter, in fact, consists of a
392 more robust and flexible algorithm, particularly suited for the classification of a large number of samples and
393 is suggested to be a more powerful tool for reliable variable selection compared to the traditional LDA (Rashid,
394 Hussain, Ahmad, & Abdullah, 2019).

395 *3.3.1. Correlation and principal component analysis*

396 Pearson's associations between all the elements were reported by plotting the correlation matrix shown in
397 Fig. 1. The main significant patterns of covariations were found among most of the REEs and, in particular,
398 between Ce and La ($r > 0.9$), and among Nd, Pr, Tb, Ho, Gd, Er, Eu, Dy, and Sm ($r > 0.8$). Similarly, bivariate

399 positive correlation regarding the patterns Na–Sr–Cd–Li–Mg–Ca ($r > 0.9$), P–K ($r > 0.9$), and Cs–Rb ($r > 0.8$)
400 were also detected. Concentrations of U were positively correlated with Ca, Mg, Pd, Li, Cd, Sr, and Na ($r >$
401 0.8) as well as amounts of Co with Cu, Mo, and V ($r > 0.7$). The most important and significant negative
402 correlations were instead found among K–Rb–Cs–Mn and among Na–Sr–Cd–Li–Ca–Mg–U ($r < -0.7$) patterns.
403 The association between the characteristic element distributions and cuttlefish origins was further investigated
404 by PCA. The first PCA computation took into consideration all the fifty-two elements and, as a result, the
405 number of original variables was set by LOOCV to six final principal components (PCs) enclosing 72.6% and
406 57.3% of the explained and predictive data variance, respectively. The first two PCs of the model contributed
407 for 27.5% and 20.8% of the total explained variance, with the PC2 enclosing the fraction of information related
408 to sample provenance, as reported in Fig. 2A. The distribution of Chioggia cuttlefish along the negative axis
409 of the PC2 was mainly dominated by the highest contribution of Rb and K, followed by Mn, Al, Cs, P, As,
410 and, at lower levels, Bi (see loading vectors reported in Fig. 2A). On the other hand, French Atlantic cuttlefish
411 distribution was mainly guided by contribution from Na, Sr, Cd, Li, Mg, Ca, U, and Mo, whose vectors angles
412 nearness underlined the strong covariation previously stated by CA. Likewise, a strong contribution and a high
413 degree of covariation was confirmed in PCA for REEs, which impacted on the negative axis of PC1.
414 During this preliminary stage, one sample from Chioggia was also found to be a strong outlier since it fell
415 outside the 95% confidence interval defined by the Hotelling's T^2 range (see Fig. 2A). For this reason, it was
416 excluded from subsequent statistics.

417 Since the most influential loading values were found for a total of 16 out 52 elements (Li, B, Na, Mg, Ca, Co,
418 Cu, Zn, Sr, Mo, Ru, Pd, Cd, Hf, Pt, and U), variable filtering was performed (see Section 2.7) and PCA was
419 thus recomputed. Loading weights (0-1 scaled) of selected elements are reported in Supplementary Materials,
420 Table S4. The reduced PCA model extracted three PCs that explained 84.5% of the data variability. The PC1
421 (explained variability, 54.1%; predictive variability, 48.3%) was mainly characterized by Zn, Cd, Pd, Ca, U,
422 Sr, while in the PC2 (explained variability, 16.3%; predictive variability, 17.1%) Pt was the most influential
423 element (Fig. 2B). Hence, maximum variation in the original dataset was retained while reducing the number
424 of significant features. At the same time, sample separation in the score space was significantly expressed
425 along the PC1 using the selected variables rather than the whole set of elements.

426 *3.3.2. Classification of cuttlefish by origin: LDA and PLS-DA approaches using selected variables*

427 The LW-LDA classification model for cuttlefish origins, created by using variables previously selected by
428 PCA (Li, B, Na, Mg, Ca, Co, Cu, Zn, Sr, Mo, Ru, Pd, Cd, Hf, Pt, and U) was described by two discriminant
429 functions (DFs) explaining 99.7% of the data variability. The DF1 (explained variability, 83.1%; canonical
430 correlation, 97.2%) and the DF 2 (explained variability, 16.6%; canonical correlation, 86.7%), whose statistical
431 significance was confirmed by low Wilk's Lambda values (0.004 and 0.13, respectively, $p = 0.000$) were
432 responsible for the clusterization of cuttlefish in the bidimensional score plot reported in Fig. 3A. Samples
433 from French Atlantic were well discriminated from samples from Chioggia and Mediterranean Sea by the DF1
434 and Chioggia samples well discriminated from Mediterranean samples by the DF2. The most important
435 loadings for the DF1 were Sr and Na, while Co was found to be the most influent element for the DF2 (Fig.
436 3A). Based on these results, 100% of cuttlefish were correctly recognized in CV, but, when performing the
437 external test set validation, two samples from Mediterranean area were misclassified as samples from French
438 Atlantic and one French Atlantic sample as of Mediterranean origin, thus resulting in an overall accuracy of
439 the model of 97% (Table 3). Fisher's discriminant function coefficients for each geographical provenance are
440 reported in Supplementary Material, Table S5.

441 In S-LDA the forward selection method based on Wilk's Lambda values was chosen to select the
442 discriminating variables, taking 3.84 as the minimum partial F value to include a variable and 2.71 as the
443 maximum F value to exclude a variable from the model (see Supplementary materials, Supplementary
444 Experimental Section). Based on this, two DFs and 12 out 52 discriminating elements were finally included
445 into the model, corresponding to Na, Co, B, K, Cd, V, U, Rb, Ni, Ba, Cu, and As (Supplementary Materials,
446 Table S6), of which Na, Co, B, Cd, U, and Cu were the same as those selected in LW-LDA.

447 The DF1 and DF2 enclosed 82.5% and 17.5% of variance of original data, with a canonical correlation of
448 97.4% and 88.5% respectively. The significant discriminatory ability of each DF was confirmed not only by
449 Wilk's Lambda values of 0.03 for the DF1 and 0.114 for the DF2 ($p = 0.000$), but also by 99% accuracy in
450 LOOCV (one Mediterranean sample erroneously classified as Atlantic sample) and 100% accuracy in external
451 validation (Table 3). Fisher's coefficients for each provenance are reported in Supplementary Materials, Table
452 S6.

453 Optimal classification results (100% sensitivity, specificity, and accuracy, Table 3) were obtained by the
454 application of PLS-DA when 21 elements were selected on the basis of their VIP scores ($VIP > 1$) and used as

455 classificatory variables (see Supplementary Materials, Table S7). Among these elements, ten were found to be
456 the same of those previously included in stepwise-LDA. Although Sr, Mn, Mo, Li, Ca, Mg, Se, Bi, Cs, P, and
457 Y were selected in addition, some of these (Sr, Mo, Li, Ca, and Mg) were shared with LW-LDA. The highest
458 VIP scores were 1.672 for V and 1.628 for Co, thus indicating the maximum contribution of these elements to
459 the geographical separation between cuttlefish.

460 The VIP-PLS-DA model was created by using six LVs explaining 90.1% of variance ($R^2X = 0.901$), of which
461 92.2% was directly correlated with labels of groups ($R^2Y = 0.922$). In addition, the overall training model's
462 predictive power of 88.8% ($Q^2 = 0.888$), resulted in 100% of samples to be correctly classified both in LOOCV
463 and external validation (Table 3), without any possible misleading interpretation deriving from overfit or
464 overprediction of the model as assessed by permutation test (average R^2Y -y-intercept = 0.048; average Q^2 -y-
465 intercept = -0.371). Further information about this validation is reported in Supplementary Materials,
466 Supplementary Fig. S4.

467 As for the RMSECV and RMSEP values, these were found to be low (0.156 and 0.159, respectively) and very
468 close to each other, thus remarking the outstanding ability of the simplified model in categorizing cuttlefish.

469 Fig. 3B shows the bidimensional score and loading graphs of the first two DFs obtained by S-LDA, while the
470 first two LVs obtained by VIP-PLS-DA are reported in Fig. 3C. In both cases, cuttlefish were well separated
471 from each other, with samples from Chioggia distributing in the lower right quadrant of the score plots. In
472 addition, both the DF1 and the LV1 were mainly responsible for sample discrimination in the score plot.

473 By looking at the loading plots, Chioggia cuttlefish were found to be positively correlated with U, V, Ni, Ba,
474 As, Rb, and K in S-LDA (Fig. 3B) and with Cs, Rb, K and P in PLS-DA (Fig. 3C). The highest loading scores
475 for Mediterranean cuttlefish were instead observed for Cd in S-LDA, which in turn was diagonally opposite
476 to the elements that characterized Chioggia samples. In PLS-DA model, the highest degree of correlation with
477 Mediterranean cuttlefish was instead found for V, Se, Co, Cu, Mo, Ni, Bi, and Mn while for French Atlantic
478 cuttlefish the highest contribution was exerted by Na together with Mg, U, Li, Sr, Ca and Cd. Likewise, Na
479 showed a high loading value on the DF2 negative axis and, therefore, it allowed to distinguish French samples
480 from the other groups in the S-LDA model.

481 According to the results of multi-element analysis of fish and seafood products already published in literature,
482 some of the most discriminant elements found in the present study might be linked to anthropogenic pollution
483 (As, V, Cd and U) or to geogenic sources (Cs and Mn) (Costas-Rodríguez et al., 2010).
484 Consistently to the results achieved, the major elements Na, Mg, and K were also previously identified among
485 the most useful indicators of geographical origin of crabs (Luo et al., 2019) and prawns (Gopi et al., 2019).
486 The variation in the amounts of Na, Mg and P in the tissues of Pacific shrimps was also correlated with specific
487 sampling sites, whose waters were characterized by higher salinity (Li, Han, Dong, & Boyd, 2019).
488 The concentrations of As which, in the present work, was one of the selected discriminating elements for
489 Chioggia cuttlefish, were also reported in mussels from Venice Lagoon (in which Chioggia is located) as being
490 positively associated to the higher degree of salinity of this area and thus valuable for the identification of local
491 products (Cubadda, Raggi, & Coni, 2006). Moreover, also Ni, Co, Se, Mo, and V amounts showed a strong
492 link with anthropic sources of element contamination of the Venice Lagoon (Cubadda et al., 2006), therefore
493 the specific pattern distribution of these trace elements in Chioggia cuttlefish tissues might be supposed to
494 specifically reflect their origin.
495 Based on these results, the use of multivariate classifiers in combination with the pre-selection of the most
496 origin-discriminant elements, was proved to be an efficient, rapid and smart analytical strategy to assure the
497 authenticity of the provenance of cuttlefish samples. Although each of the classification techniques applied
498 showed satisfying results, suggesting that possible users may choose the most convenient methodology to suit
499 the specific needs, superior and unequivocal outcomes were obtained by applying VIP-PLS-DA.

500 **4. Conclusions**

501 Cephalopods are among some of the most consumed fishery products and are appreciated by the consumers
502 all over the world but, to the best of our knowledge, scientific insights concerning their elemental profile and
503 the authentication of their geographical origin are still lacking.

504 The results presented in this study revealed for the first time that the geographical imprint of Italian traditional
505 cuttlefish from Chioggia can be extracted through the simultaneous quantification of more than fifty elements
506 by ICP-MS and successfully used to discriminate the product from cuttlefish originating from different areas.

507 The whole elemental profile was elaborated by means of different chemometric techniques. In particular, three
508 independent variable selection strategies merged with linear or regression pattern recognition multivariate
509 techniques (LW-LDA, S-LDA, and VIP-PLS-DA) were tested to develop classification rules while reducing
510 the number of variables to the lowest possible, in order to find the most parsimonious way that best described
511 sample origins. Although the application of VIP-PLS-DA led to the extraction of the highest number of
512 variables, analysis of performance metrics suggested the best results for this methodology, since values of
513 sensitivity, specificity, and classification accuracy of 100% were achieved in internal and in external validation
514 thanks to the contribution of Na, K, Ca, P, Mg, Cu, Co, Mn, Se, Ni, Mo, Li, B, Ba, Bi, Sr, As, V, Rb, Cs, Y,
515 U, and Cd.

516 In summary, the elemental pattern linked to the geographical origin of cuttlefish appears to be determined by
517 a combination of macro, trace and ultra-trace elements of both natural and anthropogenic origin, which are
518 known to be absorbed by the animals from the surrounding environment. In particular, the contribution of
519 anthropogenic elements here strongly emerges as a key analytical determinant for cephalopods authenticity
520 assessment. Of note, also concentrations of some heavy toxic metals such as Cd and As, although being so low
521 as not to represent a potential health risk, were useful for the discrimination purpose.

522 Considering the permanent and continuous release of all these elements in the aquatic environment deriving
523 from modern industrial and agricultural practices, the quantification of anthropogenic contaminants in fishery
524 products should therefore be encouraged, as well as further specific critical evaluations of their variation within
525 seawater is recommended. In such a scenario, the promising results obtained may pave the way for practical
526 application of the proposed methodology in the fishery sector to trace back different geographical origins, but
527 also for the protection and ongoing promotion of traditional local fishery products for which a quality mark
528 has been recognized.

529 **Appendix A. Supplementary Materials**

530 Supplementary data associated with this article can be found, in the online version, at

531 **Declaration of interest:** The authors declare that they have no known competing financial interests or
532 personal relationships that could have appeared to influence the work reported in this paper.

533 **Data Availability:** The dataset generated during the current study is available from the corresponding author
534 Prof. Lenka Husáková on reasonable request.

535 **Acknowledgements**

536 The authors gratefully acknowledge the financial support from the University of Pardubice (project no.
537 SGS_2020_002) and Dr. Giovanni Muresu Ibbi (DAVIMAR S.r.l, Chioggia, Italy) for providing cuttlefish
538 samples.

539 **References**

- 540 Abbas, O., Zadavec, M., Baeten, V., Mikuš, T., Lešić, T., Vulić, A., Prpić, J., Jemeršić, L., & Pleadin, J.
541 (2018). Analytical methods used for the authentication of food of animal origin. *Food Chemistry*, 246,
542 6–17. <https://doi.org/10.1016/j.foodchem.2017.11.007>
- 543 Arbuckle, N. S. M., & Wormuth, J. H. (2014). Trace elemental patterns in Humboldt squid statoliths from
544 three geographic regions. *Hydrobiologia*, 725(1), 115–123. <https://doi.org/10.1007/s10750-013-1608-4>
- 545 Ayas, D., & Ozogul, Y. (2011). The Effects of Season and Sex in the Metal Levels of Mature Common
546 Cuttlefish (*Sepia officinalis*) in Mersin Bay, Northeastern Mediterranean, *Journal of Food Science*,
547 76(4), 121–124. <https://doi.org/10.1111/j.1750-3841.2011.02152.x>
- 548 Bizzi, C. A., Pedrotti, M. F., Silva, J. S., Barin, J. S., Nóbrega, J. A., & Flores, E. M. M. (2017). Microwave-
549 assisted digestion methods: Towards greener approaches for plasma-based analytical techniques.
550 *Journal of Analytical Atomic Spectrometry*, 32(8), 1448–1466. <https://doi.org/10.1039/c7ja00108h>
- 551 Bosch, A. C., O'Neill, B., Sigge, G. O., Kerwath, S. E., & Hoffman, L. C. (2015). Heavy metals in marine
552 fish meat and consumer health: a review. *Journal of the Science of Food and Agriculture*, 96, 32–48.
553 <https://doi.org/10.1002/jsfa.7360>
- 554 Bua, G. D., Albergamo, A., Annuario, G., Zammuto, V., Costa, R., & Dugo, G. (2017). High-Throughput
555 ICP-MS and Chemometrics for Exploring the Major and Trace Element Profile of the Mediterranean
556 Sepia Ink. *Food Analytical Methods*, 10(5), 1181–1190. <https://doi.org/10.1007/s12161-016-0680-6>
- 557 Bustamante, P., Lahaye, V., Durnez, C., Churlaud, C., & Caurant, F. (2006). Total and organic Hg
558 concentrations in cephalopods from the North Eastern Atlantic waters: Influence of geographical origin
559 and feeding ecology. *Science of the Total Environment*, 368(2–3), 585–596.
560 <https://doi.org/10.1016/j.scitotenv.2006.01.038>
- 561 Carvalho, M. L., Santiago, S., & Nunes, M. L. (2005). Assessment of the essential element and heavy metal
562 content of edible fish muscle. *Analytical and Bioanalytical Chemistry*, 382, 426–432.
563 <https://doi.org/10.1007/s00216-004-3005-3>
- 564 Chaguri, M. P., Maulvault, A. L., Nunes, M. L., Santiago, D. A., Denadai, J. C., Fogaça, F. H., Sant'Ana, L.
565 S., Ducatti, C., Bandarra, N., Carvalho, M. L., Marques, A. (2015). Different tools to trace geographic

566 origin and seasonality of croaker (*Micropogonias furnieri*). *LWT - Food Science and Technology*,
567 61(1), 194–200. <https://doi.org/10.1016/j.lwt.2014.11.006>

568 Clodia database (2017). Database of Fishery Data from Chioggia, Northern Adriatic Sea.
569 <http://chioggia.biologia.unipd.it/banche-dati/>. Accessed 10.03.21

570 Commission of the European Communities (2006). Commission regulation (EC) No 1881/2006 of 19
571 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Official Journal of the*
572 *European Union*, L364/5 (20.12.2006). Retrieved from [https://eur-lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32006R1881)
573 [content/EN/ALL/?uri=CELEX%3A32006R1881](https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32006R1881)

574 Costas-Rodríguez, M., Lavilla, I., & Bendicho, C. (2010). Classification of cultivated mussels from Galicia
575 (Northwest Spain) with European Protected Designation of Origin using trace element fingerprint and
576 chemometric analysis. *Analytica Chimica Acta*, 664(2), 121–128.
577 <https://doi.org/10.1016/j.aca.2010.03.003>

578 Cubadda, F., Raggi, A., & Coni, E. (2006). Element fingerprinting of marine organisms by dynamic reaction
579 cell inductively coupled plasma mass spectrometry. *Analytical and Bioanalytical Chemistry*, 384(4),
580 887–896. <https://doi.org/10.1007/s00216-005-0256-6>

581 Danezis, G. P., Tsagkaris, A. S., Camin, F., Brusic, V., & Georgiou, C. A. (2016). Food authentication:
582 Techniques, trends & emerging approaches. *TrAC Trends in Analytical Chemistry*, 85, 123-132.
583 <https://doi.org/10.1016/j.trac.2016.02.026>

584 Drivelos, S. A., & Georgiou, C. A. (2012). Multi-element and multi-isotope-ratio analysis to determine the
585 geographical origin of foods in the European Union. *TrAC - Trends in Analytical Chemistry*, 40, 38–51.
586 <https://doi.org/10.1016/j.trac.2012.08.003>

587 European Food Safety Authority (EFSA), Panel on Contaminants in the Food Chain (CONTAM), (2009).
588 Scientific Opinion on Arsenic in Food. *EFSA Journal*, 7(10), 1351.
589 <https://doi.org/10.2903/j.efsa.2009.1351>.

590 European Parliament and Council of the European Union. (2013). Regulation (EU) No 1379/2013 of the the
591 European Parliament and of the Council of 11 December 2013 on the common organization of the
592 markets in fishery and aquaculture products, amending Council Regulations (EC) No 1184/2006 and

593 (EC) No 1224/2009 and repealin Council Regulation (EC) No 104/2000. *Official Journal of the*
594 *European Union*, L354(28.12.2013), 1–21. Retrieved from [https://eur-lex.europa.eu/legal-](https://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1588369230953&uri=CELEX:32013R1379)
595 [content/EN/TXT/?qid=1588369230953&uri=CELEX:32013R1379](https://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1588369230953&uri=CELEX:32013R1379)

596 FAO (Food and Agriculture Organization of the United Nations) (2020). FAO Species Fact Sheets, *Sepia*
597 *officinalis* (Linnaeus, 1758). <http://www.fao.org/fishery/species/2711/en>. Accessed 06.03.20

598 FAO/WHO (Food and Agriculture Organization of the United Nations/World Health Organization)
599 (2004). Evaluations of the Joint FAO/WHO Expert Committee on Food Additives (JECFA). From
600 twenty-sixth to seventy-fourth meetings. [https://apps.who.int/food-additives-contaminants-jecfa-](https://apps.who.int/food-additives-contaminants-jecfa-database/search.aspx)
601 [database/search.aspx](https://apps.who.int/food-additives-contaminants-jecfa-database/search.aspx). Accessed 03.05.20

602 Gonzalez, M. H., Souza, G. B., Oliveira, R. V., Forato, L. A., Nóbrega, J. A., & Nogueira, A. R. A. (2009).
603 Microwave-assisted digestion procedures for biological samples with diluted nitric acid: Identification
604 of reaction products. *Talanta*, 79(2), 396–401. <https://doi.org/10.1016/j.talanta.2009.04.001>

605 Gopi, K., Mazumder, D., Sammut, J., Saintilan, N., Crawford, J., & Gadd, P. (2019b). Combined use of
606 stable isotope analysis and elemental profiling to determine provenance of black tiger prawns (*Penaeus*
607 *monodon*). *Food Control*, 95, 242–248. <https://doi.org/10.1016/j.foodcont.2018.08.012>

608 Husáková, L., Urbanová, I., Šrámková, J., Černohorský, T., Krejčová, A., Bednaříková, M., Frýdová E.,
609 Nedělková, I., & Pilařová, L. (2011). Analytical capabilities of inductively coupled plasma orthogonal
610 acceleration time-of-flight mass spectrometry (ICP-oe-TOF-MS) for multi-element analysis of food and
611 beverages. *Food Chemistry*, 129(3), 1287–1296. <https://doi.org/10.1016/j.foodchem.2011.05.047>

612 Husáková, L., Urbanová, I., Šídová, T., Cahová, T., Faltys, T., & Šrámková, J. (2015). Evaluation of
613 ammonium fluoride for quantitative microwave-assisted extraction of silicon and boron from different
614 solid samples. *International Journal of Environmental Analytical Chemistry*, 95(10), 922–935.
615 <https://doi.org/10.1080/03067319.2015.1070409>

616 Iguchi, J., Isshiki, M., Takashima, Y., Yamashita, Y., & Yamashita, M. (2014). Identifying the origin of
617 *Corbicula* clams using trace element analysis. *Fisheries Science*, 80(5), 1089–1096.
618 <https://doi.org/10.1007/s12562-014-0775-1>

619 Kang, X., Zhao, Y., Shang, D., Zhai, Y., Ning, J., & Sheng, X. (2018). Elemental analysis of sea cucumber

620 from five major production sites in China: A chemometric approach. *Food Control*, 94(March), 361–
621 367. <https://doi.org/10.1016/j.foodcont.2018.07.019>

622 Lall, S.P. (2002). The minerals. J.E. Halver, R.P. Hardy (Eds.), *Fish Nutrition* (3rd ed.), Academic Press,
623 London, pp. 260–308

624 Li, L., Boyd, C. E., & Sun, Z. (2016). Authentication of fishery and aquaculture products by multi-element
625 and stable isotope analysis. *Food Chemistry*, 194, 1238–1244.
626 <https://doi.org/10.1016/j.foodchem.2015.08.123>

627 Li, L., Han, C., Dong, S., & Boyd, C. E. (2019). Use of elemental profiling and isotopic signatures to
628 differentiate Pacific white shrimp (*Litopenaeus vannamei*) from freshwater and seawater culture areas.
629 *Food Control*, 95(August 2018), 249–256. <https://doi.org/10.1016/j.foodcont.2018.08.015>

630 Li, Y., Chen, C., Li, B., Sun, J., Wang, J., Gao, Y., Zhao, Y., Chai, Z. (2006). Elimination efficiency of
631 different reagents for the memory effect of mercury using ICP-MS. *Journal of Analytical Atomic*
632 *Spectrometry*, 21(1), 94–96. <https://doi.org/10.1039/b511367a>

633 Luo, R., Jiang, T., Chen, X., Zheng, C., Liu, H., & Yang, J. (2019). Determination of geographic origin of
634 Chinese mitten crab (*Eriocheir sinensis*) using integrated stable isotope and multi-element analyses.
635 *Food Chemistry*, 274(March 2018), 1–7. <https://doi.org/10.1016/j.foodchem.2018.08.104>

636 Ministerial Decree 5 June 2014. XIV Revisione dell'Elenco Nazionale dei Prodotti Agroalimentari
637 Tradizionali. *G.U. Serie Generale n.141 del 20-06-2014 - Suppl. Ordinario n. 48*, Ministero delle
638 Politiche Agricole Alimentari e Forestali, MIPAAF: Rome, Italy, 2014. Retrieved from
639 <https://www.gazzettaufficiale.it/eli/id/2014/06/20/14A04557/sg>

640 Neff, J. M. (2002). Chapter 3 - Arsenic in the Ocean. *Bioaccumulation in Marine Organisms, Effect of*
641 *Contaminants from Oil Well Produced Water* (1st ed.). Elsevier Science, pp. 57–78.
642 <https://doi.org/10.1016/B978-008043716-3/50004-X>

643 Noack, C. W., Dzombak, D. A., & Karamalidis, A. K. (2014). Rare earth element distributions and trends in
644 natural waters with a focus on groundwater. *Environmental Science and Technology*, 48(8), 4317–
645 4326. <https://doi.org/10.1021/es4053895>

646 Ortea, I., & Gallardo, J. M. (2015). Investigation of production method, geographical origin and species

647 authentication in commercially relevant shrimps using stable isotope ratio and/or multi-element
648 analyses combined with chemometrics: An exploratory analysis. *Food Chemistry*, 170, 145–153.
649 <https://doi.org/10.1016/j.foodchem.2014.08.049>

650 Raimundo, J., Pereira, P., Vale, C., & Caetano, M. (2005). Fe , Zn , Cu and Cd concentrations in the
651 digestive gland and muscle tissues of *Octopus vulgaris* and *Sepia officinalis* from two coastal areas in
652 Portugal. *Ciencias Marinas*, 31(1B), 243–251. <https://doi.org/10.7773/cm.v31i12.91>

653 Rashid, N. A., Hussain, W. S. E. C., Ahmad, A. R., & Abdullah, F. N. (2019). Performance of classification
654 analysis: A comparative study between PLS-DA and integrating PCA+LDA. *Mathematics and*
655 *Statistics*, 7(4), 24–28. <https://doi.org/10.13189/ms.2019.070704>

656 Smith, G. (2018). Step away from stepwise. *Journal of Big Data*, 5(1). [https://doi.org/10.1186/s40537-018-](https://doi.org/10.1186/s40537-018-0143-6)
657 0143-6

658 Smith, R. G., & Watts, C. A. (2009). Determination of the country of origin of farm-raised shrimp (family
659 penaeide) using trace metal profiling and multivariate statistics. *Journal of Agricultural and Food*
660 *Chemistry*, 57(18), 8244–8249. <https://doi.org/10.1021/jf901658f>

661 Storelli, M. M., Giacomini-Stuffler, R., Storelli, A., & Marcotrigiano, G. O. (2007). Cadmium and
662 mercury in cephalopod molluscs : Estimated weekly intake. *Food Additives and Contaminants*, 23(1),
663 25–30. <https://doi.org/10.1080/02652030500242023>

664 Varrà, M. O., Ghidini, S., Zanardi, E., Badiani, A., & Ianieri, A. (2019). Authentication of European sea bass
665 according to production method and geographical origin by light stable isotope ratio and rare earth
666 elements analyses combined with chemometrics. *Italian Journal of Food Safety*, 8(1).
667 <https://doi.org/10.4081/ijfs.2019.7872>

668 **Figure Captions**

669 **Fig. 1.** Pearson's correlation heat-map of the investigated elements.

670 **Fig. 2.** Biplot resulting from PCA applied to all the elements (A) and selected elements (B). The sample
671 surrounded by the red dotted line in the biplot indicate an outlier according to Hotelling's T^2 test (95%
672 confidence interval). Blue triangles: samples from Chioggia (FAO fishing area 37.2.1); red diamonds: samples
673 from Mediterranean Sea (FAO fishing area 37.1/37.2); yellow circles: samples from French Atlantic (FAO
674 fishing area 27.7.e).

675 **Fig. 3.** Comparison of score (left) and loading graphs (right) from LW-LDA (A), S-LDA (B) and VIP-PLS-
676 DA (C) for cuttlefish samples from different origins. Blue triangles: samples from Chioggia (FAO fishing area
677 37.2.1); red diamonds: samples from Mediterranean Sea (FAO fishing area 37.1/37.2); yellow circles: samples
678 from French Atlantic (FAO fishing area 27.7.e).

679

680

681

682

683

684

685

686

Table 1

Agilent 7900 ICP-MS operating conditions.

Parameter	Setting		
ICP			
Plasma mode	General purpose		
Rf power (27 MHz) (W)	1550		
Sampling depth (mm)	8		
Plasma gas flow (L min ⁻¹)	15		
Auxiliary gas flow (L min ⁻¹)	0.9		
Nebulizer gas flow (L min ⁻¹)	1		
Nebulizer pump (rps)	0.1		
Spray chamber temperature (°C)	2		
Mass spectrometer	No gas mode	He mode	HEHe mode ^a
Extract 1 (V)		0	
Extract 2 (V)		-250	
Omega bias (V)	-100	-120	-120
Omega lens (V)	9.7	7.8	9.6
Cell entrance	-30	-40	-140
Cell exit	-50	-60	-150
Deflect (V)	11.6	1	-77
Plate bias	-35	-60	-150
Helium flow (mL min ⁻¹)	0	5	10
OctP bias	-8	-18	-100
OctP RF		200	
Energy discrimination (V)		5	
Number of elements	39 ^b	12 ^c	5 ^d
Acquisition			
Points per peak	1		
Replicates	3		
Sweeps/replicate	100		
Total acquisition time (s)	75		

^a HEHe mode - high energy helium mode; Monitored isotopes (integration time): ^b ⁷Li, ¹¹B, ²⁴Mg, ⁶⁶Zn, ⁸⁵Rb, ⁸⁸Sr, ⁸⁹Y, ⁹⁰Zr, ⁹⁵Mo, ¹⁰¹Ru, ¹⁰³Rh, ¹⁰⁵Pd, ¹¹¹Cd, ¹¹⁸Sn, ¹²¹Sb, ¹³³Cs, ¹³⁸Ba, ¹³⁹La, ¹⁴⁰Ce, ¹⁴¹Pr, ¹⁴⁶Nd, ¹⁴⁷Sm, ¹⁵³Eu, ¹⁵⁷Gd, ¹⁵⁹Tb, ¹⁶³Dy, ¹⁶⁵Ho, ¹⁶⁶Er, ¹⁷²Yb, ¹⁷⁵Lu, ¹⁷⁸Hf, ¹⁸⁵Re, ¹⁹⁵Pt, ²⁰⁵Tl, ²⁰⁶⁺²⁰⁷⁺²⁰⁸Pb, ²⁰⁹Bi, ²³²Th, ²³⁸U (all 0.1 s); ^c ²³Na (0.3 s), ²⁷Al (0.1 s), ³⁹K, ⁴⁴Ca (both 0.3 s), ⁵¹V (1 s), ⁵²Cr, ⁵⁵Mn, ⁵⁶Fe, ⁵⁹Co, ⁶⁰Ni, ⁶³Cu, ¹⁰³Rh (all 0.3 s); ^d ³¹P (0.1 s), ⁷⁵As, ⁷⁸Se (both 1 s), ¹⁰³Rh (0.3 s).


Table 2 Multi-elemental composition of cuttlefish from different geographical origins. Elements are sorted according to decreasing median concentrations of cuttlefish from Chioggia.



Element	Chioggia (n = 17*3)				Mediterranean Sea (n = 25*3)				French Atlantic (n = 26*3)			
	Mean \pm ME	Median	Min	Max	Mean \pm ME	Median	Min	Max	Mean \pm ME	Median	Min	Max
Na	11164 \pm 867	11235 ^a	8659	14187	14334 \pm 486	14290 ^b	12015	16338	24632 \pm 1332	24730 ^c	17932	30749
K	13471 \pm 1067	13768 ^a	9113	16488	10648 \pm 402	10547 ^b	8898	12400	8353 \pm 531	8252 ^c	6196	12039
P	10697 \pm 831	10635 ^a	7459	1617	9084 \pm 221	9079 ^b	7440	9883	8203 \pm 308	8176 ^c	6791	10401
Mg	2297 \pm 207	2274 ^a	1626	3180	2613 \pm 75	2643 ^a	2164	2871	3767 \pm 183	3776 ^b	2898	4795
Ca	820 \pm 102	751 ^a	619	1338	1299 \pm 53	1304 ^b	1019	1513	1781 \pm 118	1760 ^c	1239	2493
As	79 \pm 9	75.8 ^a	56.1	121.3	69 \pm 4.3	67.4 ^{ab}	50.5	98.8	60 \pm 4.7	62.3 ^b	35.5	77.1
Li	0.25 \pm 0.024	0.24 ^a	0.18	0.34	0.34 \pm 0.011	0.33 ^b	0.28	0.38	0.50 \pm 0.025	0.51 ^c	0.36	0.65
Zn	62 \pm 5.2	60.7 ^a	41.1	86.3	73 \pm 1.6	72.8 ^b	63.8	80.7	74 \pm 3.5	73.8 ^b	58.6	94.1
Al	16 \pm 5.5	10.7 ^a	4.7	43.8	7 \pm 1.9	5.69 ^b	2.97	22.7	7 \pm 2.5	5.44 ^b	2.87	34.4
Cu	10 \pm 1.8	8.55 ^a	5.93	20.2	24 \pm 3.4	21.3 ^b	12.5	42.9	20 \pm 3.4	16.3 ^b	8.29	36.2
Sr	9 \pm 1.1	8.83 ^a	6.69	14.9	14.8 \pm 0.44	14.9 ^b	12.0	16.8	25 \pm 1.6	26.0 ^c	16.3	33.2
Fe	11 \pm 4	7.85 ^a	3.62	33.0	9 \pm 1.9	8.19 ^a	4.32	25.0	8 \pm 2	5.59 ^a	3.41	21.7
B	6.3 \pm 0.56	6.06 ^{ab}	4.46	8.20	5.3 \pm 0.29	5.35 ^a	4.11	7.15	7.6 \pm 0.85	7.88 ^b	4.15	11.8
Rb	5.3 \pm 0.42	5.17 ^a	3.94	6.99	4.5 \pm 0.14	4.61 ^a	3.93	5.13	3.6 \pm 0.22	3.57 ^b	2.56	5.27
Se	1.3 \pm 0.17	1.33 ^a	0.881	2.11	2.0 \pm 0.14	1.91 ^b	1.47	2.69	1.65 \pm 0.083	1.62 ^c	1.26	2.16
Mn	2.2 \pm 0.87	1.28 ^a	0.93	6.86	2.5 \pm 0.52	2.04 ^a	1.41	6.59	1.0 \pm 0.23	0.83 ^b	0.54	3.57
Ni	0.4 \pm 0.52	0.36 ^a	0.22	0.76	0.52 \pm 0.056	0.50 ^b	0.27	1.98	0.28 \pm 0.034	0.26 ^a	0.15	0.48
Cr	0.18 \pm 0.060	0.18 ^a	0.037	0.55	0.15 \pm 0.041	0.13 ^a	0.068	0.54	0.22 \pm 0.082	0.13 ^a	0.081	1.07
Ba	0.3 \pm 0.13	0.18 ^a	0.087	1.17	0.22 \pm 0.031	0.16 ^a	0.12	0.50	0.21 \pm 0.039	0.18 ^a	0.11	0.43
Pb	0.10 \pm 0.024	0.093 ^a	0.050	0.23	0.10 \pm 0.019	0.099 ^a	0.063	0.31	0.09 \pm 0.017	0.046 ^a	0.077	0.20
V	0.07 \pm 0.036	0.040 ^a	0.018	0.26	0.25 \pm 0.040	0.22 ^b	0.12	0.47	0.11 \pm 0.023	0.11 ^a	0.039	0.27
Zr	0.04 \pm 0.018	0.027 ^a	0.0059	0.13	0.05 \pm 0.015	0.036 ^{ab}	0.0071	0.14	0.2 \pm 0.20	0.051 ^b	0.0066	2.55
Co	0.04 \pm 0.019	0.025 ^a	0.016	0.15	0.21 \pm 0.023	0.21 ^b	0.11	0.31	0.14 \pm 0.026	0.14 ^c	0.058	0.31
Mo	0.029 \pm 0.0066	0.024 ^a	0.018	0.060	0.10 \pm 0.013	0.089 ^b	0.061	0.18	0.10 \pm 0.019	0.084 ^b	0.044	0.23
Pd	0.016 \pm 0.0013	0.016 ^a	0.012	0.020	0.024 \pm 0.0054	0.021 ^a	0.012	0.067	0.029 \pm 0.0033	0.027 ^b	0.017	0.059
Cs	0.017 \pm 0.0015	0.016 ^a	0.012	0.023	0.0146 \pm 0.00085	0.014 ^a	0.012	0.023	0.012 \pm 0.0014	0.012 ^b	0.0081	0.027
Y	0.02 \pm 0.012	0.015 ^a	0.0057	0.11	0.008 \pm 0.0012	0.0074 ^b	0.0039	0.017	0.01 \pm 0.013	0.0077 ^b	0.0050	0.17
Bi	0.178 \pm 0.0022	0.017 ^a	0.011	0.027	0.023 \pm 0.0040	0.021 ^a	0.015	0.065	0.010 \pm 0.0041	0.0066 ^b	0.0029	0.044
Sn	0.015 \pm 0.0071	0.011 ^a	0.0044	0.061	0.022 \pm 0.0049	0.019 ^b	0.0075	0.051	0.015 \pm 0.0040	0.011 ^a	0.0052	0.053
Cd	0.02 \pm 0.017	0.0099 ^a	0.0035	0.13	0.08 \pm 0.013	0.071 ^b	0.035	0.15	0.25 \pm 0.059	0.22 ^c	0.067	0.58
Sb	0.007 \pm 0.0028	0.0053 ^a	0.0023	0.021	0.009 \pm 0.0012	0.0084 ^b	0.0046	0.016	0.0075 \pm 0.00080	0.0068 ^b	0.0052	0.014
U	0.0032 \pm 0.00076	0.0026 ^a	0.0019	0.0074	0.0043 \pm 0.00042	0.0040 ^a	0.0027	0.0069	0.010 \pm 0.0015	0.0091 ^b	0.0049	0.019
Th	0.002 \pm 0.0011	0.0016 ^a	0.00081	0.0082	0.005 \pm 0.0036	0.0022 ^a	0.0011	0.046	0.003 \pm 0.0019	0.0015 ^a	0.0011	0.024
Ce*	29 \pm 10	20.6 ^a	3.56	82.0	44 \pm 20	24.0 ^a	11.4	250	26 \pm 8.9	18.8 ^a	8.31	91.7
La*	20 \pm 11	11.9 ^{ab}	2.70	90.87	29 \pm 12	20.3 ^a	8.03	150	16 \pm 5.6	10.5 ^b	5.09	56.0
Nd*	8 \pm 4.7	3.98 ^a	1.58	33.2	5 \pm 1.5	4.45 ^a	1.74	20.1	4 \pm 1.0	2.82 ^b	1.33	12.0
Hf*	2.5 \pm 0.52	2.23 ^a	1.15	4.94	8 \pm 8.4	2.89 ^{ab}	1.42	103	8 \pm 4.9	4.19 ^b	0.92	57.3
Tl*	1.5 \pm 0.11	1.42 ^a	1.07	1.94	1.4 \pm 0.51	1.21 ^b	0.83	7.30	1.3 \pm 0.61	1.05 ^b	0.62	8.66
Pr*	2 \pm 1.0	0.96 ^a	0.38	7.43	1.2 \pm 0.34	0.97 ^a	0.45	4.62	0.8 \pm 0.23	0.63 ^b	0.33	2.56
Sm*	1.5 \pm 0.81	0.74 ^a	0.25	5.04	0.9 \pm 0.28	0.82 ^a	0.34	3.73	0.7 \pm 0.18	0.55 ^a	0.33	2.32
Dy*	1.1 \pm 0.44	0.74 ^a	0.34	3.22	0.8 \pm 0.18	0.66 ^{ab}	0.27	2.21	0.7 \pm 0.17	0.51 ^b	0.33	2.02
Gd*	1 \pm 0.5	0.69 ^{ab}	0.24	3.20	0.8 \pm 0.19	0.72 ^a	0.32	2.52	0.6 \pm 0.15	0.50 ^b	0.31	1.97
Er*	0.7 \pm 0.23	0.54 ^a	0.20	1.73	0.5 \pm 0.10	0.42 ^{ab}	0.23	1.17	0.4 \pm 0.11	0.39 ^b	0.19	1.31
Yb*	0.7 \pm 0.21	0.51 ^a	0.22	1.48	0.39 \pm 0.078	0.37 ^{ab}	0.14	0.93	0.4 \pm 0.10	0.34 ^b	0.16	1.31
Eu*	0.4 \pm 0.18	0.24 ^{ab}	0.12	1.19	0.28 \pm 0.061	0.25 ^a	0.15	0.87	0.22 \pm 0.044	0.19 ^b	0.12	0.62
Ho*	0.22 \pm 0.078	0.17 ^a	0.062	0.60	0.15 \pm 0.030	0.14 ^{ab}	0.047	0.39	0.14 \pm 0.037	0.11 ^b	0.063	0.45
Tb*	0.20 \pm 0.082	0.14 ^a	0.056	0.58	0.15 \pm 0.037	0.12 ^{ab}	0.055	0.43	0.12 \pm 0.030	0.095 ^b	0.054	0.37
Ru*	0.10 \pm 0.029	0.087 ^a	0.031	0.24	0.14 \pm 0.027	0.13 ^a	0.054	0.34	0.14 \pm 0.023	0.13 ^a	0.044	0.29
Re*	0.10 \pm 0.032	0.081 ^{ab}	0.016	0.24	0.12 \pm 0.019	0.11 ^a	0.070	0.24	0.09 \pm 0.023	0.072 ^b	0.025	0.28
Lu*	0.09 \pm 0.032	0.074 ^a	0.023	0.23	0.06 \pm 0.015	0.047 ^a	0.021	0.18	0.06 \pm 0.017	0.053 ^a	0.021	0.22
Pt*	0.13 \pm 0.034	0.10 ^a	0.060	0.25	0.2 \pm 0.12	0.12 ^a	0.062	1.53	0.15 \pm 0.061	0.10 ^a	0.019	0.60
Hg [#]	0.20 \pm 0.030	0.18 ^a	0.15	0.34	0.20 \pm 0.023	0.17 ^a	0.14	0.38	0.21 \pm 0.018	0.20 ^a	0.13	0.29


Concentrations are expressed as mg kg⁻¹ (d.m.) and are reported as mean values \pm margin of error (ME) at 95% confidence level. Min–Max: minimum and maximum values found. *: concentrations are expressed as μ g kg⁻¹. Hg[#]: determined by direct mercury analyzer AMA254. Values followed by different superscript letters (a–c) in the same row are significantly different ($p \leq 0.05$).

Table 3 Supervised classification model performances in cross-validation (CV) and in external test set validation.

Model	Validation	Overall classification rate (%)	Sensitivity (%)	Specificity (%)	Accuracy (%)
LW-LDA	CV	100	100	100	100
	External	96	96	98	97
S-LDA	CV	99	99	99	99
	External	100	100	100	100
VIP-PLS-DA	CV	100	100	100	100
	External	100	100	100	100







Supplementary Materials

Multi-element signature of cuttlefish and its potential for the discrimination of different geographical provenances and traceability

Maria Olga VARRÀ^a, Lenka HUSÁKOVÁ^{b}, Jan PATOČKA^b, Sergio*

GHIDINI^a, Emanuela ZANARDI^{a}*

^aDepartment of Food and Drug, University of Parma, Parma, Via del Taglio, 10, Parma 43126, Italy

^bDepartment of Analytical Chemistry, Faculty of Chemical Technology, University of Pardubice, Studentska 573 HB/D, Pardubice, CZ-532 10, Czech Republic. Tel. +420 466 037 029; Fax: +420 466 037 068.

E-mail address: lenka.husakova@upce.cz

Table of Contents

Supplementary Experimental Section	1
Figure S1. Internal standard (ISTD) stability for sequence including 111 samples of cuttlefish.	6
Table S1. Principal spectral interferences encountered on the Agilent 7900	7
Table S2. Comparison of measured and certified concentrations in selected control standards.....	8
Figure S2. Graphical summary of recovery values of the different certified reference materials analyzed .	13
Table S3. Method Detection Limits (MDL) and Method Limits of Quantification (MLOQ) ($\mu\text{g kg}^{-1}$) and normalized calibration slopes (NCS) ($1/\mu\text{g L}^{-1}$) for the analysis of different elements in cuttlefish samples.	14
Figure S3. Graphical summary of Method Detection Limits (MDL) and Method Limits of Quantification (MLOQ)	15
Table S4. Loading weights from PCA performed using all the elements	16
Table S5. Fisher's classification coefficients for the LW-LDA model.....	17
Table S6. Variables selected according to Wilk's Lambda values used for the creation of stepwise-LDA models and relative Fisher's classification coefficients	18
Table S7. VIP values used for the creation of VIP-PLS-DA model and relative regression coefficients	19
Figure S4. Permutation plot (400 random permutations) assessing the validity of VIP-PLS-DA training model.....	20

Supplementary Experimental Section

Essential elements such as calcium, iron, copper, and zinc are frequently normally distributed in fish tissues since concentrations are metabolically regulated by the organisms. Contrarily, non-essential trace elements' concentrations are completely dependent on the exposure to the specific environment and, therefore, frequently characterized by a positive skewed lognormal distribution (Phillips & Russo, 1978). As expected, both data homoscedasticity and normality assumptions tested by the Shapiro-Wilk's and Levene's tests were violated for the largest portion of the elements in the analyzed cuttlefish samples ($p \leq 0.05$), thus nonparametric statistics was used to investigate and describe elemental concentrations, taking into consideration that, alongside with this, the main advantage of nonparametric statistical tests is the low sensitivity to small samples sizes and to the potential bias of outlying data (Helsel & Hirsch, 1992). The comparison among groups of cuttlefish was therefore reported after having performed the nonparametric Kruskal–Wallis analysis of variance followed Dunn's post hoc (Sawilowsky & Fahoome 2014) test for multiple comparison ($p \leq 0.05$).

The first chemometric technique applied to analyze the huge amount of ICP-MS data obtained from cuttlefish sample was principal component analysis (PCA). The PCA is an unsupervised projection method aimed at dimensionality reduction of the original set of correlated variables into new fewer uncorrelated latent variables extracting as much systematic variation as possible of the original data. PCA is mostly used as a preliminary explorative tool to study any intrinsic correlation among observations, variables as well as the bidirectional relation between variables and observation and it is a powerful technique for the rapid detection of strong multivariate outliers (Jolliffe & Cadima, 2016). In the present work, the application of PCA to elemental data was mainly aimed at obtaining the first global description of dataset, after having corrected the non-normal data distribution by Box-Cox transformation (Sakia, 1992) and the wide range of magnitude of values by Z-score standardization (Shiffler, 1988). In combination with Pearson's correlation analysis applied to Box-Cox transformed data, PCA was also useful for data collinearity reduction necessary for the subsequent step in supervised classification.

To define the optimal number of principal components to retain and avoid data overfitting and overoptimistic results, leave-one-out cross validation (LOOCV) following the approach of Krzanowski

was employed (Eastment & Krzanowski, 1982). According to this, one observation per time was retained and predicted by the model which, in turn, was trained on the remaining observations. This sequence was repeated until all observations were kept out one by one. Thus, considering the sample size, LOOCV statistics resulted from a number of 68*3 trained models. The selected internal validation method was chosen taking into consideration the effective number of independent samples in the dataset and because since it is faster to be computed compared to other CV methods.

In study dealing with elemental fingerprints, linear discriminant analysis (LDA) is the most frequently used supervised classification method. By using Euclidean distance, LDA is based on defining new linear combinations of the original variables able to maximize the separation between class of samples while minimizing intra-class variability. The maximum likelihood ratio criterion, referred to as Wilks' lambda ($p \leq 0.05$) was applied to verify the statistical significance of each discriminant function, where smaller values suggested greater discriminatory power associated to the function (Todorov, 2007). Since LDA may be prone to failing in classification, especially when the samples size for each class is greatly unbalanced, equal priors in estimations were not set, but probabilities were calculated based on unequal sample sizes (Sanchez, 1974; Xue & Titterington, 2008).

With the aim to define a classification model by using the minimum number of variables that increased between-group variability and, at the same time, decreased within-class variability, forward stepwise-LDA (S-LDA) was applied to Box-Cox transformed and Z-score standardized data. An *F*-statistic was used to define the statistical significance of each variable to discrimination, where *F*-to-remove value (indicating the cut off value for a variable to be excluded) was set to 2.71 and *F*-to-enter value (indicating the cut-off value for a variable to be included) was set to 3.84.

Standard LDA and S-LDA were evaluate in terms of recognition ability in LOOCV applied to 66% of original observations (training set), i.e. the percentage of samples correctly assigned to the proper class. Considering that the internal cross-validation is often insufficient to accurately assess the predictability of the model, an external validation of the training model was also performed using the excluded 33% of data which were randomly but consistently selected from the whole dataset. External test observations which do not uniformly cover the range of training set distribution, may led to misleading results (Consonni, Ballabio & Todeschini, 2010).

Partial least square-discriminant analysis (PLS-DA) is a regression-based supervised technique aimed at finding interrelation between original variables and categorical variables by building new (latent) variables for the maximum separation between the different groups of samples. In particular, the categorical variable matrix is transformed into a dummy variable matrix, which boundaries of classification of samples into one class are defined using Bayes theorem. In the present work, the quality and of discriminant models based on PLS-DA of Box-Cox transformed and Z-score standardized data was assessed by applying LOOCV on the 66% of observation of the training set (Eastment & Krzanowski, 1982) and by evaluating the resulting estimating of fitting (R^2X and R^2Y) and predictive ability (Q^2) and the root-mean square error of cross-validation (RMSECV) (Bellabio & Consonni, 2013). Here too, the external validation of the regression model was performed using the independent 33% of observation left out during the calibration step (test set) and the number of samples correctly classified was evaluated. The reliability, in terms overoptimistic fitting and predictability results, was further checked by permutation tests (400 random permutation). The significance of the model was thus confirmed if the y-intercept values of the R^2Y were ≤ 0.4 and y-intercept values of Q^2 were ≤ 0.05 (Van der Voet, 1994). When building classification regression models, the discrimination among informative, redundant, and noisy variables is often an important step. The variable-influence on projection (VIP) index was used to identify and select the most discriminant variables in PLS-DA (Andersen & Bro, 2010), which were afterward employed to build a new simplified model. The VIP index summarizes the cumulative importance of each variable and represents the weighted sum of squares of the PLS weights, considering the whole explained variability related to responses (sample groups) in all extracted components (Andersen & Bro, 2010). VIP indexes higher than one are considered to be the most significant for explaining the correlation of the observation to all the responses (Andersen & Bro, 2010). The overall quality and robustness of all the supervised models was assessed by taking into consideration the experimental percentages of true positive samples (sensitivity %), true negative samples (specificity %) and prediction accuracy (%). These metrics were calculated as follows, following what has been previously reported by Fawcett (2006):

$$\text{Sensitivity (\%)} = \frac{\text{TP}}{\text{TP} + \text{FN}} * 100$$

$$\text{Specificity (\%)} = \frac{\text{TN}}{\text{TN} + \text{FP}} * 100$$

$$\text{Accuracy (\%)} = \frac{\text{TN} + \text{TP}}{\text{TN} + \text{TP} + \text{FN} + \text{FP}} * 100$$

Whereas: TP = true positive samples

TN = true negative samples

FP = false positive samples

FN = false negative samples

LIST OF REFERENCES

- Andersen, C. M., & Bro, R. (2010). Variable selection in regression—a tutorial. *Journal of Chemometrics*, 24(11- 12), 728-737. <https://doi.org/10.1002/cem.1360>
- Ballabio, D., & Consonni, V. (2013). Classification tools in chemistry. Part 1: linear models. PLS-DA. *Analytical Methods*, 5(16), 3790-3798. <https://doi.org/10.1039/C3AY40582F>
- Berrueta, L. A., Alonso-Salces, R. M., & Héberger, K. (2007). Supervised pattern recognition in food analysis. *Journal of chromatography A*, 1158(1-2), 196-214. <https://doi.org/10.1016/j.chroma.2007.05.024>
- Consonni, V., Ballabio, D., & Todeschini, R. (2010). Evaluation of model predictive ability by external validation techniques. *Journal of chemometrics*, 24(3- 4), 194-201. <https://doi.org/10.1002/cem.1290>
- Downey, G. (Ed.). (2016). *Advances in food authenticity testing*. Woodhead Publishing
- Eastment, H. T., & Krzanowski, W. J. (1982). Cross-Validatory Choice of the Number of Components from a Principal Component Analysis. *Technometrics*, 24(1), 73-77. <https://doi.org/10.1080/00401706.1982.10487712>

- Fawcett, T. (2006). An introduction to ROC analysis. *Pattern Recognition Letters*, 27(8), 861–874. <https://doi.org/10.1016/j.patrec.2005.10.010>
- Helsel and Hirsch 1992 *Statistical methods in water resources*: Amsterdam the Netherlands, Elsevier Studies in Environmental Science, 49, 522.
- Jolliffe, I., & Cadima, J. (2016). Principal component analysis: A review and recent developments. *Philosophical Transactions A*, 374(2065), 20150202. <https://doi.org/10.1098/rsta.2015.0202>
- Phillips, G. R., & Russo, R. C. (1978). Metal bioaccumulation in fishes and aquatic invertebrates: a literature review. U.S. Environmental Protection Agency, Washington, D.C., EPA/600/3-78/103.
- Sakia, R. M. (1992). The Box- Cox transformation technique: a review. *Journal of the Royal Statistical Society: Series D (The Statistician)*, 41(2), 169-178. <https://doi.org/10.2307/2348250>
- Shiffler, R. E. (1988). Maximum Z scores and outliers. *The American Statistician*, 42(1), 79-80. <https://doi.org/10.1080/00031305.1988.10475530>
- Sanchez, P. M. (1974). The unequal group size problem in discriminant analysis. *Journal of the Academy of Marketing Science*, 2(4), 629–633.
- Todorov, V. (2007). Robust selection of variables in linear discriminant analysis. *Statistical Methods and Applications*, 15, 395–407 <https://doi.org/10.1007/s10260-006-0032-6>
- Van der Voet, H. (1994). Comparing the predictive accuracy of models using a simple randomization test. *Chemometrics and Intelligent Laboratory Systems*, 25, 313-323. [https://doi.org/10.1016/0169-7439\(94\)85050-X](https://doi.org/10.1016/0169-7439(94)85050-X).
- Xue, J. H., & Titterton, D. M. (2008). Do unbalanced data have a negative effect on LDA? *Pattern Recognition*, 41(5), 1558-1571. <https://doi.org/10.1016/j.patcog.2007.11.008>
- Sawilowsky, S., & Fahoome, G. (2014). Kruskal- Wallis Test: Basic. Wiley StatsRef: Statistics Reference Online. <https://doi.org/10.1002/9781118445112.stat06567>

Figure S1. Internal standard (ISTD) stability for sequence including 111 samples of cuttlefish (sample name is displayed on the x-axis) measured during the 8-hour run. Due to limited space, not all sample names are shown in the X-axis labels. The ISTD recoveries are displayed relative to the calibration blank.




Table S1. Principal spectral interferences encountered on the Agilent 7900 from the major components of food matrices, plasma gas and sample solvent onto the analysis of selected elements.

Interfered isotope	Interfering ion
$^{27}\text{Al}^+$	$^{12}\text{C}^{15}\text{N}^+$, $^{12}\text{C}^{14}\text{NH}^+$
$^{31}\text{P}^+$	$^{14}\text{N}^{16}\text{OH}^+$, $^{15}\text{N}^{16}\text{O}^+$,
$^{39}\text{K}^+$	$^{23}\text{Na}^{16}\text{O}^+$, $^{38}\text{ArH}^+$
$^{44}\text{Ca}^+$	$^{12}\text{C}^{16}\text{O}^{16}\text{O}^+$
$^{48}\text{Ca}^+$	$^{31}\text{P}^{17}\text{O}^+$, $^{31}\text{P}^{16}\text{O}^1\text{H}^+$
$^{51}\text{V}^+$	$^{35}\text{Cl}^{16}\text{O}^+$, $^{37}\text{Cl}^{14}\text{N}^+$
$^{52}\text{Cr}^+$	$^{40}\text{Ar}^{12}\text{C}^+$, $^{35}\text{Cl}^{16}\text{OH}^+$, $^{37}\text{Cl}^{14}\text{NH}^+$
$^{53}\text{Cr}^+$	$^{40}\text{Ar}^{13}\text{C}^+$, $^{35}\text{Cl}^{18}\text{O}^+$, $^{37}\text{Cl}^{16}\text{O}^+$, $^{36}\text{Ar}^{16}\text{OH}^+$
$^{55}\text{Mn}^+$	$^{39}\text{K}^{16}\text{O}^+$, $^{37}\text{Cl}^{18}\text{O}^+$, $^{23}\text{Na}^{32}\text{S}^+$
$^{54}\text{Fe}^+$	$^{37}\text{Cl}^{17}\text{O}^+$, $^{37}\text{Cl}^{16}\text{O}^1\text{H}^+$
$^{56}\text{Fe}^+$	$^{40}\text{Ar}^{16}\text{O}^+$, $^{40}\text{Ca}^{16}\text{O}^+$
$^{57}\text{Fe}^+$	$^{40}\text{Ar}^{16}\text{OH}^+$, $^{40}\text{Ca}^{16}\text{OH}^+$
$^{58}\text{Ni}^+$	$^{40}\text{Ca}^{18}\text{O}^+$, $^{40}\text{Ar}^{18}\text{O}^+$, $^{23}\text{Na}^{35}\text{Cl}^+$
$^{59}\text{Co}^+$	$^{43}\text{Ca}^{16}\text{O}^+$, $^{42}\text{Ca}^{16}\text{O}^1\text{H}^+$, $^{40}\text{Ar}^{18}\text{OH}^+$
$^{60}\text{Ni}^+$	$^{44}\text{Ca}^{16}\text{O}^1+$, $^{43}\text{Ca}^{16}\text{O}^1\text{H}^+$, $^{23}\text{Na}^{37}\text{Cl}^+$
$^{61}\text{Ni}^+$	$^{44}\text{Ca}^{16}\text{O}^1\text{H}^+$, $^{36}\text{Ar}^{25}\text{Mg}^+$, $^{38}\text{Ar}^{23}\text{Na}^+$, $^{23}\text{Na}^{37}\text{ClH}^+$
$^{62}\text{Ni}^+$	$^{23}\text{Na}^{23}\text{Na}^{16}\text{O}^+$, $^{44}\text{Ca}^{18}\text{O}^+$, $^{36}\text{Ar}^{26}\text{Mg}^+$
$^{63}\text{Cu}^+$	$^{40}\text{Ar}^{23}\text{Na}^+$, $^{31}\text{P}^{16}\text{O}^{16}\text{O}^+$, $^{12}\text{C}^{16}\text{O}^{35}\text{Cl}^+$, $^{12}\text{C}^{14}\text{N}^{37}\text{Cl}^+$
$^{64}\text{Zn}^+$	$^{31}\text{P}^{16}\text{O}^{17}\text{O}^+$, $^{32}\text{S}^{16}\text{O}_2^+$, $^{32}\text{S}_2^+$, $^{36}\text{Ar}^{12}\text{C}^{16}\text{O}^+$, $^{38}\text{Ar}^{12}\text{C}^{14}\text{N}^+$, $^{48}\text{Ca}^{16}\text{O}^+$
$^{65}\text{Cu}^+$	$^{31}\text{P}^{16}\text{O}^{18}\text{O}^+$, $^{32}\text{S}^{33}\text{S}^+$, $^{33}\text{S}^{16}\text{O}_2^+$, $^{32}\text{S}^{16}\text{O}_2^1\text{H}^+$, $^{32}\text{S}_2\text{H}^+$, $^{14}\text{N}^{16}\text{O}^{35}\text{Cl}^+$, $^{48}\text{Ca}^{16}\text{O}^+$
$^{66}\text{Zn}^+$	$^{34}\text{S}^{16}\text{O}_2^+$, $^{32}\text{S}^{16}\text{O}^{18}\text{O}^+$, $^{32}\text{S}^{17}\text{O}_2^+$, $^{33}\text{S}^{16}\text{O}^{17}\text{O}^+$, $^{32}\text{S}^{34}\text{S}^+$, $^{33}\text{S}_2^+$, $^{48}\text{Ca}^{18}\text{O}^+$
$^{67}\text{Zn}^+$	$^{33}\text{S}^{34}\text{S}^+$, $^{34}\text{S}^{16}\text{O}^{17}\text{O}^+$, $^{33}\text{S}^{16}\text{O}^{18}\text{O}^+$, $^{32}\text{S}^{17}\text{O}^{18}\text{O}^+$, $^{33}\text{S}^{17}\text{O}_2^+$, $^{33}\text{S}_2\text{H}^+$, $^{48}\text{Ca}^{18}\text{OH}^+$, $^{14}\text{N}^{16}\text{O}^{37}\text{Cl}^+$, $^{16}\text{O}_2^{35}\text{Cl}^+$
$^{68}\text{Zn}^+$	$^{34}\text{S}_2^+$, $^{32}\text{S}^{18}\text{O}_2^+$
$^{69}\text{Ga}^+$	$^{34}\text{S}_2\text{H}^+$, $^{32}\text{S}^{18}\text{O}_2\text{H}^+$, $^{16}\text{O}_2^{37}\text{Cl}^+$
$^{71}\text{Ga}^+$	$^{34}\text{S}^{18}\text{O}_2\text{H}^+$
$^{72}\text{Ge}^+$	$^{40}\text{Ar}^{32}\text{S}^+$, $^{35}\text{Cl}^{37}\text{Cl}^+$, $^{40}\text{Ar}^{16}\text{O}_2^+$
$^{73}\text{Ge}^+$	$^{40}\text{Ar}^{33}\text{S}^+$, $^{35}\text{Cl}^{37}\text{ClH}^+$, $^{40}\text{Ar}^{16}\text{O}_2\text{H}^+$
$^{74}\text{Ge}^+$	$^{40}\text{Ar}^{33}\text{S}^+$, $^{37}\text{Cl}_2^+$, $^{39}\text{K}^{35}\text{Cl}^+$
$^{75}\text{As}^+$	$^{43}\text{Ca}^{16}\text{O}_2^+$, $^{40}\text{Ca}^{35}\text{Cl}^+$, $^{40}\text{Ar}^{34}\text{SH}^+$
$^{77}\text{Se}^+$	$^{40}\text{Ca}^{37}\text{Cl}^+$, $^{40}\text{Ar}^{37}\text{Cl}^+$
$^{78}\text{Se}^+$	$^{41}\text{K}^{37}\text{Cl}^+$, $^{38}\text{Ar}^{40}\text{Ca}^+$
$^{80}\text{Se}^+$	$^{40}\text{Ar}_2^+$, $^{40}\text{Ca}_2^+$, $^{40}\text{Ar}^{40}\text{Ca}^+$
$^{82}\text{Se}^+$	$^{32}\text{S}^{17}\text{O}_2^{16}\text{O}^+$, $^{33}\text{S}^{16}\text{O}_2^{17}\text{O}^+$

Table S2. Comparison of measured and certified concentrations in selected control standards, recoveries (R), and intra-day and inter-day relative standard deviation (RSD).

Analyte	Reference sample	Declared (mg kg ⁻¹)	Found ^a (mg kg ⁻¹)	R ^b (%)	RSD (%)	
					Intra	Inter
⁷ Li	NCS ZC73015 Milk Powder	0.040	0.0414 ± 0.0032	104	3.86	6.70
	CRM 12-2-04 Wheat bread flour	0.020	0.0195 ± 0.0021	98	5.45	6.07
¹¹ B	NCS ZC73015 Milk Powder	1.56 ± 0.22	1.53 ± 0.08	98	2.75	4.85
	CRM 12-2-04 Wheat bread flour	< 0.5	0.49 ± 0.04	^d	3.96	7.08
	CRM12-2-03 Lucerne	30	30.2 ± 2.0	101	3.35	9.20
²³ Na	BCR 184 Bovine muscle	2000	2080 ± 56	104	1.35	2.07
	BCR-CRM 185 Bovine Liver	2100	2101 ± 72	102.80	1.71	3.32
	NIST 1577 Bovine Liver	2033 ± 64	1918 ± 88	94	2.31	3.78
	NIST 1566 Oyster Tissue	5100 ± 3000	5013 ± 250	98	2.48	3.02
	NCS ZC 73015 Milk Powder	4700 ± 300	4400 ± 752	94	8.57	6.79
	CRM12-2-03 Lucerne	474 ± 23	474 ± 24	100	2.51	3.53
²⁴ Mg	CRM 12-2-01 Bovine Liver	650 ± 112	645 ± 60	99	4.68	5.54
	BCR 184 Bovine muscle	1020	1026 ± 30	101	1.39	6.38
	NIST 1577 Bovine Liver	620 ± 42	594 ± 16	96	1.30	1.40
	NIST 1566 Oyster Tissue	1280 ± 90	1316 ± 78	103	2.95	8.28
	NCS ZC 73015 Milk Powder	960 ± 70	880 ± 38	92	2.24	5.77
	CRM 12-2-04 Wheat bread flour	556 ± 29	554 ± 7.2	99	0.65	7.85
	CRM12-2-03 Lucerne	3520 ± 125	3475 ± 336	99	4.82	6.44
²⁷ Al	CRM 12-2-04 Wheat bread flour	3	3.4 ± 0.2	113	1.39	3.16
	CRM12-2-03 Lucerne	330	330 ± 5	100	0.78	9.20
³¹ P	BCR 184 Bovine muscle	8300	8351 ± 328	101	1.97	2.62
	BCR-CRM 185 Bovine Liver	11700	11618 ± 452	99	1.94	11.2
	NIST 1566 Oyster Tissue	8100	8131 ± 324	101	1.99	4.91
	NCS ZC 73015 Milk Powder	7600 ± 300	7300 ± 448	96	3.17	5.30
	CRM 12-2-04 Wheat bread flour	2280 ± 85	2289 ± 95	100	2.07	2.24
	CRM12-2-03 Lucerne	3030 ± 90	2982 ± 102	98	1.71	4.60
³⁹ K	CRM 12-2-04 Wheat bread flour	2550 ± 100	2596 ± 83	102	1.60	3.73
	BCR 184 Bovine muscle	16600	16354 ± 200	99	0.61	2.57
	BCR-CRM 185 Bovine Liver	11200	10807 ± 350	97	1.62	3.60
	NIST 1577 Bovine Liver	10230 ± 640	10270 ± 512	101	2.49	4.01
	NIST 1566 Oyster Tissue	9690 ± 50	9610 ± 514	99	2.67	3.83
	NCS ZC 73015 Milk Powder	12500 ± 500	11910 ± 312	95	1.33	2.74
⁴⁴ Ca	CRM12-2-03 Lucerne	18700 ± 650	18897 ± 638	101	1.69	1.66
	BCR 184 Bovine muscle	150	151 ± 4	101	1.17	3.54
	BCR-CRM 185R Bovine Liver	131	133 ± 5	102	1.87	3.73
	NIST 1566 Oyster Tissue	1500 ± 200	1400 ± 14	93	0.49	2.28
	NCS ZC 73015 Milk Powder	9400 ± 300	8630 ± 740	92	4.29	4.75
	CRM12-2-03 Lucerne	17500 ± 750	17348 ± 229	99	0.65	1.30

Table S2. Continued

Analyte	Reference sample	Declared (mg kg ⁻¹)	Found ^a (mg kg ⁻¹)	R ^b (%)	RSD (%)	
					Intra	Inter
⁵¹ V	NIST 1566 Oyster Tissue	2.3 ± 0.1	2.1 ± 0.1	91	1.43	8.27
	CRM 12-2-01 Bovine Liver	0.26 ± 0.05	0.27 ± 0.01	104	1.73	3.14
	NCS ZC 73015 Milk Powder	0.06	0.050 ± 0.002	83	1.93	3.03
⁵² Cr	BCR 184 Bovine muscle	0.076	0.075 ± 0.003	99	2.13	7.28
	CRM 12-2-01 Bovine Liver	0.044	0.048 ± 0.002	109	2.25	9.74
	NCS ZC 73015 Milk Powder	0.39 ± 0.04	0.40 ± 0.02	103	2.22	1.98
	CRM12-2-03 Lucerne	0.900	0.74 ± 0.03	82	2.02	7.00
⁵⁵ Mn	CRM 12-2-04 Wheat bread flour	22.6 ± 1.1	22.4 ± 0.9	99	1.93	8.13
	BCR 184 Bovine muscle	0.334 ± 0.028	0.320 ± 0.017	96	2.68	4.36
	BCR-CRM 185R Bovine Liver	9.3 ± 0.3	9.4 ± 0.2	101	0.90	10.9
	NIST 1577 Bovine Liver	10.46 ± 0.47	10.2 ± 0.4	98	2.15	2.53
	NIST 1566 Oyster Tissue	17.5 ± 1.2	16.4 ± 2.3	94	7.02	6.73
	CRM 12-2-01 Bovine Liver	7.6 ± 0.5	7.46 ± 0.09	98	0.58	4.20
	NCS ZC 73015 Milk Powder	0.51 ± 0.17	0.50 ± 0.08	98	2.85	9.56
	CRM12-2-03 Lucerne	34.2 ± 1.15	34.7 ± 0.3	101	0.44	1.86
⁵⁶ Fe	BCR 184 Bovine muscle	79 ± 2	78 ± 3	99	1.80	2.72
	BCR-CRM 185R Bovine Liver	214 ± 5	217 ± 6	101	1.40	13.8
	NIST 1577 Bovine Liver	495 ± 32.5	501 ± 24	101	2.39	3.15
	NIST 1566 Oyster Tissue	195 ± 34	183 ± 18	94	4.81	3.62
	CRM 12-2-01 Bovine Liver	495 ± 28	501 ± 24	101	2.39	3.15
	NCS ZC 73015 Milk Powder	7.8 ± 1.3	6.8 ± 0.4	87	2.83	3.06
	CRM 12-2-04 Wheat bread flour	23.8 ± 1.5	22.7 ± 1.3	95	2.97	3.00
	CRM12-2-03 Lucerne	355 ± 18	350 ± 36	99	5.08	1.68
⁵⁹ Co	NIST 1577 Bovine Liver	0.300 ± 0.018	0.303 ± 0.003	101	0.48	1.44
	NIST 1566 Oyster Tissue	0.400	0.320 ± 0.028	80	4.65	7.39
	CRM 12-2-01 Bovine Liver	0.37 ± 0.03	0.36 ± 0.02	97	2.71	1.37
	CRM12-2-03 Lucerne	0.193 ± 0.0185	0.175 ± 0.003	91	0.97	4.79
⁶⁰ Ni	BCR 184 Bovine muscle	0.270	0.274 ± 0.018	102	3.28	9.96
	NIST 1566 Oyster Tissue	1.03 ± 0.19	0.94 ± 0.09	91	4.91	6.41
	CRM 12-2-04 Wheat bread flour	0.3	0.29 ± 0.05	97	3.22	5.70
	CRM12-2-03 Lucerne	2.54 ± 0.18	2.9 ± 0.2	114	2.60	5.89
⁶³ Cu	BCR 184 Bovine muscle	2.36 ± 0.06 ^c	2.30 ± 0.10	98	2.31	4.33
	BCR-CRM 185R Bovine Liver	189 ± 4	189 ± 3	100	0.85	9.91
	NIST 1577 Bovine Liver	275.2 ± 4.6	271 ± 8	98	1.42	1.74
	NIST 1566 Oyster Tissue	63.0 ± 3.5	62 ± 2	98	1.68	0.17
	CRM 12-2-01 Bovine Liver	26.3 ± 1.6	25.7 ± 1.4	98	2.69	4.80
	NCS ZC 73015 Milk Powder	0.51 ± 0.13	0.49 ± 0.01	96	1.33	7.66
	CRM 12-2-04 Wheat bread flour	2.77 ± 0.03	2.74 ± 0.02	99	0.36	3.49
	CRM12-2-03 Lucerne	11.7 ± 0.75	11.3 ± 0.3	97	1.11	4.87

Table S2. Continued

Analyte	Reference sample	Declared (mg kg ⁻¹)	Found ^a (mg kg ⁻¹)	R ^b (%)	RSD (%)	
					Intra	Inter
⁶⁶ Zn	BCR 184 Bovine muscle	166 ± 3	168 ± 4	101	1.34	1.45
	BCR-CRM 185 Bovine Liver	142 ± 3	140 ± 35	99	0.58	10.6
	NIST 1577 Bovine Liver	181.1 ± 1.0	180 ± 5	99	1.27	2.95
	NIST 1566 Oyster Tissue	852 ± 14	851 ± 2	99	1.18	0.93
	CRM 12-2-01 Bovine Liver	162 ± 6	162 ± 10	100	3.10	2.44
	NCS ZC 73015 Milk Powder	34 ± 2	34 ± 4	100	5.95	2.49
⁷⁵ As	BCR-CRM 185 Bovine Liver	0.024 ± 0.003	0.025 ± 0.002	104	3.45	6.77
	NIST 1566 Oyster Tissue	13.4 ± 1.9	12.9 ± 0.4	96	1.46	1.58
	CRM 12-2-01 Bovine Liver	0.110 ± 0.016	0.107 ± 0.005	97	2.42	6.20
	CRM 12-2-04 Wheat bread flour	0.017 ± 0.0046	0.0178 ± 0.001	105	2.78	0.45
	CRM12-2-03 Lucerne	0.262 ± 0.020	0.264 ± 0.017	101	3.17	2.38
⁷⁸ Se	BCR 184 Bovine muscle	0.183 ± 0.012	0.187 ± 0.014	102	3.79	3.52
	BCR-CRM 185R Bovine Liver	0.446 ± 0.013	0.444 ± 0.040	99	4.50	5.42
	NIST 1577 Bovine Liver	2.031 ± 0.045	2.03 ± 0.06	100	1.39	5.20
	NIST 1566 Oyster Tissue	2.1 ± 0.5	2.2 ± 0.5	105	2.55	9.27
	CRM 12-2-01 Bovine Liver	0.325 ± 0.014	0.33 ± 0.03	102	3.88	10.2
	CRM 12-2-04 Wheat bread flour	0.040	0.039 ± 0.005	98	6.83	5.69
	NCS ZC 73015 Milk Powder	0.11 ± 0.03	0.104 ± 0.006	95	2.75	7.36
	CRM12-2-03 Lucerne	0.050	0.051 ± 0.002	102	0.23	2.83
⁸⁵ Rb	NIST 1566 Oyster Tissue	4.45 ± 0.09	4.3 ± 0.3	97	2.85	3.81
	CRM 12-2-01 Bovine Liver	16.0 ± 2.7	16.1 ± 1.0	101	3.19	3.00
	NCS ZC 73015 Milk Powder	11.6 ± 0.7	11 ± 1	95	5.33	6.71
	CRM 12-2-04 Wheat bread flour	1.5	1.48 ± 0.08	99	2.61	2.04
	CRM12-2-03 Lucerne	16.1 ± 2.2	15.5 ± 0.8	96	2.49	5.11
⁸⁸ Sr	NIST 1566 Oyster Tissue	10.36 ± 0.56	9.91 ± 0.05	96	0.27	2.90
	NCS ZC 73015 Milk Powder	5.3 ± 0.6	4.7 ± 0.1	89	1.15	1.65
	CRM 12-2-04 Wheat bread flour	1.53 ± 0.16	1.56 ± 0.11	102	2.04	1.14
⁸⁹ Y	NCS ZC 73015 Milk Powder	8 ± 3 ^c	8.9 ± 0.4 ^c	111	1.93	3.03
⁹⁵ Mo	NIST 1577 Bovine Liver	3.30 ± 0.13	3.40 ± 0.15	103	2.18	2.46
	NIST 1566 Oyster Tissue	0.2	0.20 ± 0.02	100	4.00	5.04
	CRM 12-2-01 Bovine Liver	3.5 ± 0.6	3.8 ± 0.3	109	3.35	5.96
	NCS ZC 73015 Milk Powder	0.28 ± 0.03	0.27 ± 0.03	96	5.43	4.16
	CRM 12-2-04 Wheat bread flour	0.2	0.203 ± 0.011	102	2.80	5.16
	CRM12-2-03 Lucerne	0.200	0.191 ± 0.011	96	2.94	3.50
¹¹¹ Cd	BCR 184 Bovine muscle	0.013 ± 0.002	0.0132 ± 0.002	102	7.81	9.32
	BCR-CRM 185 Bovine Liver	0.298 ± 0.025	0.278 ± 0.008	93	1.38	1.34
	NIST 1577 Bovine Liver	0.097 ± 0.0014	0.096 ± 0.005	99	2.46	2.52
	NIST 1566 Oyster Tissue	3.5 ± 0.4	3.24 ± 0.09	93	1.42	2.00
	CRM 12-2-01 Bovine Liver	0.48 ± 0.03	0.47 ± 0.03	98	3.13	3.68
	CRM 12-2-04 Wheat bread flour	0.0415 ± 0.0032	0.039 ± 0.0003	94	0.35	1.41
	CRM12-2-03 Lucerne	0.136 ± 0.0065	0.136 ± 0.005	100	2.00	2.27

Table S2. Continued

Analyte	Reference sample	Declared (mg kg ⁻¹)	Found ^a (mg kg ⁻¹)	R ^b (%)	RSD (%)	
					Intra	Inter
¹¹⁸ Sn	CRM 12-2-04 Wheat bread flour	< 3	0.31 ± 0.04	^d	5.78	4.47
¹²¹ Sb	CRM 12-2-01 Bovine liver	0.033	0.033 ± 0.006	100	9.76	6.99
	NCS ZC 73015 Milk Powder	6 ^c	5.98 ± 0.26 ^c	99	2.23	11.6
	CRM 12-2-03 Lucerne	0.080	0.075 ± 0.008	94	5.12	4.00
¹³³ Cs	NIST 1577 Bovine Liver	21.7 ± 1.4 ^c	22 ± 3 ^c	101	6.91	12.1
	CRM 12-02-01 Bovine liver	0.047	0.044 ± 0.001	94	1.12	3.88
	NCS ZC 73015 Milk Powder	0.034 ± 0.005	0.031 ± 0.005	91	7.54	10.1
	CRM 12-2-04 Wheat bread flour	5 ^c	4.9 ± 0.8 ^c	98	8.59	14.6
	CRM 12-2-03 Lucerne	0.090	0.086 ± 0.002	96	1.37	1.75
¹³⁸ Ba	NCS ZC 73015 Milk Powder	1.0 ± 0.3	0.86 ± 0.03	86	1.77	8.89
	CRM 12-2-04 Wheat bread flour	1.5	1.58 ± 0.22	105	7.26	0.15
	CRM 12-2-03 Lucerne	23.4 ± 2.1	23.6 ± 0.7	101	1.57	4.06
¹³⁹ La	CRM 12-2-01 Bovine Liver	0.070	0.071 ± 0.004	101	2.66	7.36
	CRM12-2-03 Lucerne	0.940	0.95 ± 0.03	101	1.73	4.00
¹⁴⁰ Ce	CRM12-2-03 Lucerne	1.0	1.04 ± 0.06	104	3.01	4.06
¹⁴¹ Pr	NCS ZC 73015 Milk Powder	0.7 ^c	0.59 ± 0.02 ^c	85	1.35	12.0
¹⁴⁶ Nd	NCS ZC 73015 Milk Powder	2 ^c	2.2 ± 0.4 ^c	110	8.82	12.7
¹⁴⁷ Sm	NCS ZC 73015 Milk Powder	0.5 ^c	0.46 ± 0.09 ^c	92	11.4	13.6
	CRM12-2-03 Lucerne	0.140	0.141 ± 0.006	101	2.16	3.27
¹⁵³ Eu	NCS ZC 73015 Milk Powder	0.4 ^c	0.45 ± 0.09 ^c	112	2.28	8.98
	CRM12-2-03 Lucerne	0.030	0.031 ± 0.001	103	1.53	2.39
¹⁵⁹ Tb	NCS ZC 73015 Milk Powder	0.7 ^c	0.61 ± 0.07 ^c	87	5.70	12.1
	CRM12-2-03 Lucerne	0.020	0.018 ± 0.001	90	2.93	3.63
¹⁶³ Dy	NCS ZC 73015 Milk Powder	0.45 ^c	0.40 ± 0.03 ^c	89	4.49	10.9
	CRM12-2-03 Lucerne	0.090	0.087 ± 0.007	97	3.89	4.96
¹⁶⁵ Ho	NCS ZC 73015 Milk Powder	0.07 ^c	0.073 ± 0.004 ^c	104	2.46	11.8
¹⁶⁶ Er	NCS ZC 73015 Milk Powder	0.16 ^c	0.19 ± 0.004 ^c	118	0.84	1.10
¹⁷⁵ Lu	CRM12-2-03 Lucerne	5 ^c	4 ± 0.2 ^c	80	3.22	6.83
¹⁷⁸ Hf	CRM12-2-03 Lucerne	0.100	0.082 ± 0.002	82	1.77	5.83
²⁰² Hg	BCR 185 Bovine Liver	0.044 ± 0.003	0.0460 ± 0.0005	105	0.54	^d
	NIST 1577c Bovine Liver	5.36 ± 0.17 ^c	5.0 ± 0.4 ^c	93	3.80	7.06
	NIST 1566 Oyster Tissue	0.057 ± 0.015	0.0528 ± 0.0004	93	0.38	^d
	CRM 12-2-01 Bovine Liver	0.37 ± 0.02	0.35 ± 0.02	95	3.14	5.92
	NCS ZC 73015 Milk Powder	2.2 ^c	2.0 ± 0.2	91	4.50	7.73
²⁰⁵ Tl	NIST 1566 Oyster Tissue	5 ^c	4.92 ± 0.05 ^c	98	1.04	10.9
	NCS ZC 73015 Milk Powder	0.9 ^c	0.89 ± 0.03 ^c	99	1.66	6.40

Table S2. Continued

Analyte	Reference sample	Declared (mg kg ⁻¹)	Found ^a (mg kg ⁻¹)	R ^b (%)	RSD (%)	
					Intra	Inter
Pb ^c	CRM 12-2-04 Wheat bread flour	0.041 ± 0.0078	0.043 ± 0.0076	104	9.97	8.64
	BCR 184 Bovine muscle	0.239 ± 0.011	0.2343 ± 0.0096	98	2.04	2.35
	BCR-CRM 185 Bovine Liver	0.501 ± 0.027	0.512 ± 0.072 ^c	102	7.03	5.92
	NIST 1566 Oyster Tissue	0.480 ± 0.040	0.475 ± 0.040	99	4.20	4.95
	CRM 12-2-01 Bovine Liver	0.71 ± 0.08	0.71 ± 0.05	100	3.79	4.54
	NCS ZC 73015 Milk Powder	0.07 ± 0.02	0.071 ± 0.003	101	1.77	6.55
	CRM12-2-03 Lucerne	1.84 ± 0.17	1.89 ± 0.07	103	1.74	4.86
²⁰⁹ Bi	NCS ZC 73015 Milk Powder	1.2 ^c	1.27 ± 0.22 ^c	106	8.89	13.8
²³² Th	NCS ZC 73015 Milk Powder	2.8 ^c	2.64 ± 0.05 ^c	99	1.00	10.5
	CRM12-2-03 Lucerne	0.110	0.109 ± 0.006	99	2.78	3.68
²³⁸ U	NIST 1566 Oyster Tissue	0.116 ± 0.006	0.112 ± 0.006	97	2.79	7.58
	NCS ZC 73015 Milk Powder	3 ^c	3.12 ± 0.14 ^c	104	2.18	5.18

^a Mean ± 2 S.D. (n = 3).

^b Recovery (%) = (Found value/Declared value)×100.

^c µg kg⁻¹

^d Not determined.

^e Pb was measured as the sum of the three most abundant isotopes, ²⁰⁶Pb⁺, ²⁰⁷Pb⁺ and ²⁰⁸Pb⁺.




Figure S2. Graphical summary of the recovery values of the eight different certified reference materials (CRMs) used to validate the analytical methods for the quantification of different elements in cuttlefish samples. B and Sn were not determined in CRM 12-02-04 wheat bread flour, therefore missing values are present in the relative plot.

Table S3. Method Detection Limits (MDL)^a and Method Limits of Quantification (MLOQ)^a ($\mu\text{g kg}^{-1}$) and normalized calibration slopes (NCS) ($1/\mu\text{g L}^{-1}$) of Agilent 7900 Q-ICP-MS for analysis of different elements in cuttlefish samples with the use of Rh as internal standard.

Analyte	Cell mode	NCS	MDL	MLOQ	Analyte	NCS	Cell mode	MDL	MLOQ
⁷ Li ⁺	No gas	5.2×10^{-3}	0.59	1.98	¹¹⁸ Sn ⁺	1.73×10^{-2}	No gas	0.49	1.6
¹¹ B ⁺	No gas	1.4×10^{-3}	5.7	19	¹²¹ Sb ⁺	2.3×10^{-2}	No gas	0.15	0.49
²³ Na ⁺	He	1.2×10^{-3}	3500	11667	¹³³ Cs ⁺	6.4×10^{-2}	No gas	0.015	0.051
²⁴ Mg ⁺	No gas	1.3×10^{-2}	15	51	¹³⁸ Ba ⁺	5.6×10^{-2}	No gas	0.55	1.83
²⁷ Al ⁺	He	1.1×10^{-4}	8.6	29	¹³⁹ La ⁺	6.4×10^{-2}	No gas	0.013	0.043
³¹ P ⁺	HE He	3.4×10^{-5}	161	536	¹⁴⁰ Ce ⁺	6.4×10^{-2}	No gas	1.50	5.0
³⁹ K ⁺	He	4.0×10^{-4}	735	2448	¹⁴¹ Pr ⁺	7.8×10^{-2}	No gas	0.010	0.033
⁴⁴ Ca ⁺	He	1.9×10^{-5}	1291	4304	¹⁴⁶ Nd ⁺	1.2×10^{-2}	No gas	0.07	0.22
⁵¹ V ⁺	He	8.1×10^{-3}	0.06	0.21	¹⁴⁷ Sm ⁺	1.1×10^{-2}	No gas	0.27	0.88
⁵² Cr ⁺	He	1.1×10^{-2}	0.97	3.2	¹⁵³ Eu ⁺	4.0×10^{-2}	No gas	0.019	0.063
⁵⁵ Mn ⁺	He	4.3×10^{-3}	2.4	8.1	¹⁵⁷ Gd ⁺	1.8×10^{-2}	No gas	0.046	0.153
⁵⁶ Fe ⁺	He	7.7×10^{-3}	6.5	22	¹⁵⁹ Tb ⁺	8.1×10^{-2}	No gas	0.0095	0.032
⁵⁹ Co ⁺	He	2.1×10^{-2}	0.12	0.41	¹⁶³ Dy ⁺	1.9×10^{-2}	No gas	0.070	0.23
⁶⁰ Ni ⁺	He	5.7×10^{-3}	6.3	21	¹⁶⁵ Ho ⁺	7.7×10^{-2}	No gas	0.054	0.18
⁶³ Cu ⁺	He	1.7×10^{-2}	2.0	6.7	¹⁶⁶ Er ⁺	2.6×10^{-2}	No gas	0.031	0.102
⁶⁶ Zn ⁺	No gas	5.8×10^{-3}	159	528	¹⁷² Yb ⁺	1.7×10^{-2}	No gas	0.047	0.157
⁷⁵ As ⁺	HE He	2.5×10^{-3}	0.49	1.6	¹⁷⁵ Lu ⁺	7.2×10^{-3}	No gas	0.011	0.035
⁷⁸ Se ⁺	HE He	3.7×10^{-4}	1.6	5.5	¹⁷⁸ Hf ⁺	2.1×10^{-3}	No gas	0.0023	0.0075
⁸⁵ Rb ⁺	No gas	4.3×10^{-2}	0.09	0.31	¹⁸⁵ Re ⁺	2.3×10^{-3}	No gas	0.12	0.39
⁸⁸ Sr ⁺	No gas	5.6×10^{-2}	0.19	0.62	¹⁹⁵ Pt ⁺	1.6×10^{-3}	No gas	0.18	0.59
⁸⁹ Y ⁺	No gas	6.8×10^{-2}	0.03	0.10	²⁰⁵ Tl ⁺	4.1×10^{-2}	No gas	0.027	0.088
⁹⁰ Zr ⁺	No gas	3.5×10^{-2}	0.11	0.35	Pb ^b	5.3×10^{-2}	No gas	0.12	0.40
⁹⁵ Mo ⁺	No gas	1.0×10^{-2}	0.58	1.93	²⁰⁹ Bi ⁺	4.4×10^{-2}	No gas	0.033	0.11
¹⁰¹ Ru ⁺	No gas	1.2×10^{-2}	0.07	0.22	²³² Th ⁺	4.7×10^{-2}	No gas	0.033	0.11
¹⁰⁵ Pd ⁺	No gas	1.34×10^{-2}	0.14	0.45	²³⁸ U ⁺	4.8×10^{-2}	No gas	0.0005	0.0017
¹¹¹ Cd ⁺	No gas	6.0×10^{-3}	0.0038	0.013	Hg ^c	2.8×10^{-2}		0.2	0.7

^a Values were calculated assuming a sample mass of 0.100 g.

^b Pb is measured as the sum of the three most abundant isotopes, ²⁰⁶Pb⁺, ²⁰⁷Pb⁺ and ²⁰⁸Pb⁺.

^c Values were evaluated for direct analysis of Hg by single purpose atomic absorption spectrometer AMA 254.




Figure S3. Graphical summary of Method Detection Limits (MDL) ($\mu\text{g kg}^{-1}$) and Method Limits of Quantification (MLOQ) ($\mu\text{g kg}^{-1}$) for the analysis of different elements in cuttlefish samples.

Table S4. Loading weights (0-1 scaled) on the first two principal components from PCA performed using all the elements. Variables with values higher than 0.7 were selected and used for the creation of LW-LDA model.

Variable	Loading Weight		Variable	Loading Weight	
	PC 1	PC 2		PC 1	PC 2
Li	0.9161	0.9828	Sn	0.4052	0.5677
B	0.6713	0.7207	Sb	0.4487	0.6867
Na	0.9835	0.9617	Cs	0.3746	0.1279
Mg	0.8714	0.9447	Ba	0.3775	0.6344
Al	0.2004	0.2489	La	0.3005	0.4644
P	0.4261	0.1216	Ce	0.2928	0.4470
K	0.4533	0.0000	Pr	0.0005	0.3591
Ca	0.8132	0.9881	Nd	0.0000	0.3737
V	0.3980	0.6934	Sm	0.0758	0.3760
Cr	0.5316	0.5639	Eu	0.0588	0.4170
Fe	0.1041	0.4436	Gd	0.0242	0.4294
Mn	0.1649	0.2719	Tb	0.0233	0.4141
Ni	0.3848	0.3796	Dy	0.0654	0.3997
Co	0.6354	0.8007	Ho	0.0248	0.4180
Cu	0.5961	0.7946	Er	0.0443	0.4210
Zn	0.4682	0.8007	Yb	0.1320	0.4139
As	0.4623	0.1563	Lu	0.1959	0.4488
Se	0.5489	0.6970	Hf	0.7509	0.6634
Rb	0.4229	0.0328	Re	0.6194	0.4359
Sr	0.9352	1.0000	Pt	0.7033	0.5132
Y	0.2576	0.3511	Tl	0.4505	0.2269
Zr	0.4817	0.6681	Pb	0.2908	0.5203
Mo	0.6908	0.9030	Bi	0.4470	0.2509
Ru	0.7650	0.5617	Th	0.2504	0.5523
Pd	0.8914	0.8030	U	0.7600	0.9832
Cd	0.9222	0.9777	Hg	0.6800	0.4359

Table S5. Fisher's classification coefficients for the LW-LDA model.

Variable selected	Fisher's classification coefficients		
	CH	MED	AT
Li	-13.496	3.080	5.994
B	14.378	0.615	-10.118
Na	-24.484	-6.308	23.663
Mg	30.145	1.805	-21.952
Ca	19.229	11.045	-24.197
Co	1.275	9.925	-9.745
Cu	3.603	2.396	-4.396
Zn	-0.484	-0.790	1.294
Sr	-49.228	-14.310	45.787
Mo	-3.046	1.769	0.438
Ru	0.305	-0.666	0.142
Pd	-0.615	1.774	-0.946
Cd	-6.960	-11.474	13.922
Hf	1.132	-0.357	-0.287
Pt	0.882	-0.344	-0.211
U	-2.835	-3.589	5.752
(Constant)	-31.893	-8.675	-24.855

CH = Cuttlefish from Chioggia (FAO 37.2.1). MED = cuttlefish from Mediterranean Sea (FAO 37.1/37.2) AT = cuttlefish from French Atlantic Ocean (FAO 27.7.e).

Table S6. Variables selected according to Wilk’s Lambda values used for the creation of stepwise-LDA models and relative Fisher’s classification coefficients for each group.

Variable selected	Lambda value	<i>T</i>	<i>F</i> to remove	<i>p</i>	Fisher’s classification coefficients		
					CH	MED	AT
					Na	0.011	0.199
Co	0.004	0.219	12.568	0.000	0.879	7.071	−7.012
B	0.006	0.207	65.093	0.000	15.871	0.960	−11.211
K	0.005	0.110	34.483	0.000	24.739	2.006	−17.469
Cd	0.004	0.171	24.886	0.000	−6.304	−12.504	14.739
V	0.005	0.319	29.741	0.000	−3.456	6.218	−3.797
U	0.004	0.194	14.543	0.000	1.876	−7.844	6.759
Rb	0.003	0.138	5.703	0.000	−9.350	0.011	5.943
Ni	0.003	0.740	7.558	0.000	1.404	2.285	−2.878
Ba	0.003	0.700	4.316	0.000	2.614	−0.112	−1.444
Cu	0.003	0.274	5.639	0.000	1.087	3.725	−3.948
As	0.003	0.428	4.393	0.000	−2.660	−1.828	3.321
(Constant)					−33.155	−10.750	−27.163

T = tolerance. *F* to remove = maximum *F* set to 2.71. *p* = significant level of 0.05. CH = Cuttlefish from Chioggia (FAO 37.2.1). MED = cuttlefish from Mediterranean Sea (FAO 37.1/37.2) AT = cuttlefish from French Atlantic Ocean (FAO 27.7.e)

Table S7. VIP values used for the creation of VIP-PLS-DA model and relative regression coefficients for each group.

Variable selected	VIP value	Regression coefficients		
		CH	MED	AT
V	1.672	-0.2584	0.4376	-0.2115
Co	1.628	-0.1722	0.2502	-0.0995
Na	1.477	-0.1700	0.0678	0.0809
Sr	1.447	-0.2113	0.1065	0.0783
Mn	1.420	0.1365	0.0292	-0.1487
Mo	1.416	-0.0867	0.0393	0.0365
Cd	1.411	0.1963	-0.4848	0.3131
Li	1.391	-0.2127	0.7827	0.0257
U	1.386	0.2916	-0.5152	0.2601
Cu	1.375	-0.0285	0.0930	-0.0681
Ni	1.370	0.0056	0.0572	-0.0620
Ca	1.368	-0.2666	0.2128	0.0204
Mg	1.361	-0.0263	-0.1046	0.1276
B	1.342	0.3117	-0.1665	-0.1062
Se	1.273	-0.1543	0.1802	-0.0451
K	1.242	0.1371	-0.0803	-0.0397
Bi	1.227	0.3117	-0.1665	-0.1062
Rb	1.174	-0.0167	0.1343	-0.1197
Cs	1.060	0.0269	-0.0675	0.0440
P	1.046	0.1007	-0.1280	0.0398
Y	1.003	0.0509	0.1523	0.0212
(Constant)		0.5546	0.7827	0.7827

CH = Cuttlefish from Chioggia (FAO 37.2.1). MED = cuttlefish from Mediterranean Sea (FAO 37.1/37.2) AT = cuttlefish from French Atlantic Ocean (FAO 27.7.e).




Figure S4. Permutation plot (400 random permutations) assessing the validity of VIP-PLS-DA training model. The intercept to the y-axis of the regression line, correlating the R2Y and Q2 values between the original and the permuted y-variables (displayed on the X-axis) and the cumulative value of R2Y and Q2 values (displayed on the X-axis), outline the degree of model's overfitting.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

CRedit authorship contribution statement

Maria Olga Varra: Writing – original draft, Investigation, Data curation, Formal analysis, Validation, Resources. **Lenka Husáková:** Writing - review & editing, Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data curation, Supervision, Funding acquisition. **Jan Patočka:** Methodology, Investigation, Validation, Formal analysis. **Emanuela Zanardi:** Conceptualization, Writing - review & editing, Project administration, Funding acquisition, Supervision. **Sergio Ghidini:** Resources, Conceptualization.